

Original Paper

Evaluating Self-Management Behaviors of Diabetic Patients in a Telehealthcare Program: Longitudinal Study Over 18 Months

Lichin Chen^{1*}; Lee-Ming Chuang^{2*}; Chia-Hsiun Chang²; Chiou-Shiang Wang³; I-Ching Wang³; Yufang Chung⁴; Hui-Yu Peng⁵; Hui-Chuen Chen⁵; Yu-Ling Hsu³; Yu-Sheng Lin¹; Huang-Jen Chen⁶; Tieng-Chun Chang²; Yi-Der Jiang²; Hung-Chang Lee⁷; Ching-Ting Tan⁸; Hsin-Lu Chang⁹; Feipei Lai^{1,6,10}

¹Graduate Institute of Biomedical Electronics and Bioinformatics, National Taiwan University, Taipei, Taiwan

²Department of Internal Medicine, National Taiwan University Hospital, National Taiwan University, Taipei, Taiwan

³Department of Nursing, National Taiwan University Hospital, National Taiwan University, Taipei, Taiwan

⁴Department of Electrical Engineering, Tunghai University, Taichung, Taiwan

⁵Department of Dietetics, National Taiwan University Hospital, National Taiwan University, Taipei, Taiwan

⁶Department of Computer Science and Information Engineering, National Taiwan University, Taipei, Taiwan

⁷Department of Information Management, Tamkang University, Taipei, Taiwan

⁸Department of Otolaryngology, College of Medicine, National Taiwan University, Taipei, Taiwan

⁹Department of Management Information Systems, National Chengchi University, Taipei, Taiwan

¹⁰Department of Electrical Engineering, National Taiwan University, Taipei, Taiwan

*these authors contributed equally

Corresponding Author:

Lichin Chen

Graduate Institute of Biomedical Electronics and Bioinformatics

National Taiwan University

No. 1, Sec 4, Roosevelt Road, Taipei, 10617, Taiwan

Taipei, 10617

Taiwan

Phone: 886 933846944

Fax: 886 233663754

Email: d98945012@ntu.edu.tw

Abstract

Background: Self-management is an important skill for patients with diabetes, and it involves frequent monitoring of glucose levels and behavior modification. Techniques to enhance the behavior changes of diabetic patients have been developed, such as diabetes self-management education and telehealthcare. Although the patients are engaged in self-management activities, barriers to behavior changes remain and additional work is necessary to address the impact of electronic media and telehealthcare on patient self-care behaviors.

Objective: The aims of this study were to (1) explore the behaviors of diabetic patients interacting with online applications, (2) determine the impact of a telehealthcare program among 7 self-care behaviors of the patients, and (3) determine the changes in glycosylated hemoglobin (HbA_{1c}) levels.

Methods: A telehealthcare program was conducted to assist the patients with 7 self-care activities. The telehealthcare program lasted for 18 months and included the use of a third-generation mobile telecommunications glucometer, an online diabetes self-management system, and a teleconsultant service. We analyzed the data of 59 patients who participated in the telehealthcare program and 103 who did not. The behavioral assessments and the HbA_{1c} data were collected and statistically analyzed to determine whether the telehealthcare services had an impact on the patients. We divided the 18-month period into 3 6-month intervals and analyzed the parameters of patients assisted by the telehealthcare service at different time points. We also compared the results of those who were assisted by the telehealthcare service with those who were not.

Results: There was a significant difference in monitoring blood glucose between the beginning and the end of the patient participation ($P=.046$) and between the overall period and the end of patient participation ($P<.001$). Five behaviors were significantly different between the intervention and control patients: being active ($P<.001$), healthy eating ($P<.001$), taking medication ($P<.001$),

healthy coping ($P=.02$), and problem solving ($P<.001$). Monitoring of blood glucose was significantly different ($P=.02$) during the 6-12 month stage of patient participation between the intervention and control patients. A significant difference between the beginning and the 6-12 month stage of patient participation was observed for the mean value of HbA_{1c} level ($P=.02$), and the differences between the overall HbA_{1c} variability and the variability of each 6-month interval was also significant.

Conclusions: Telehealthcare had a positive effect on diabetic patients. This study had enhanced blood glucose monitoring, and the patients in the program showed improvements in glycemic control. The self-care behaviors affect patient outcomes, and the changes of behavior require time to show the effects.

(*J Med Internet Res* 2013;15(12):e266) doi: [10.2196/jmir.2699](https://doi.org/10.2196/jmir.2699)

KEYWORDS

Internet; diabetes mellitus; telemedicine; self-care; online systems; personal health record; patient access to records

Introduction

Background

Self-management is an important skill for patients with diabetes mellitus [1-3], involving frequent monitoring of glucose levels and behavior modification. The primary goal of self-management is to monitor glucose metabolism and induce behavioral changes to achieve better glycemic control [3,4]. However, some patients do not respond to behavior modification and some do not have sufficient knowledge to perform self-management [5-9]. Several techniques to enhance self-care behaviors for diabetic patients have been developed, such as diabetes self-management education (DSME) and telehealthcare. DSME provides patients with self-management skills, supports behavioral changes, and achieves optimal patient outcomes in diabetes care [10-13]. Telehealthcare programs promote preventive care, self-management, and clinical consultations from a distance [6,7,9,14-16]. However, self-management is challenging because it requires behavioral changes and can be easily neglected owing to a busy lifestyle and lack of support. Barriers remain for patients who carry out self-care tasks. Previous studies have also highlighted the need for additional research to address the impact of telehealthcare and electronic media on patient self-care behaviors [17,18].

In addition to routine DSME, patients in the present study completed a telehealthcare program using a third-generation mobile telecommunication (3G) glucometer, an online diabetes self-management system, and a teleconsultant service. The aim of the present study was to assess how diabetic patients used the online diabetes self-management system, determine the impact of a telehealthcare program among the 7 self-care behaviors, and measure the changes in glycosylated hemoglobin (HbA_{1c}) level. The 7 self-care behaviors are based on the definition of the American Association of Diabetes Educators 7 Self-Care Behaviors (AADE7).

Diabetes Care in Taiwan

As effective management for chronic diseases has been established, the National Health Insurance (NHI) system of Taiwan has acknowledged the value of disease screening and disease management. The NHI has broadened payment for several disease-management programs and provided financial incentives for regular diabetes follow-up visits [19-23]. In 2001, a diabetes pay-for-performance program, known as the diabetes shared care network, was implemented [24-26]. It emphasizes

the value of a multidisciplinary care team to provide DSME and regular screening to enhance self-management skills and the early detection of complications [13].

Patients are required to return for a regular visit every 3 months for laboratory monitoring and DSME courses. Certified diabetes educators (CDEs) educate patients and evaluate their self-care behaviors and skills, which is documented in their behavioral assessments. Using past medical records and laboratory results, the CDEs attempt to determine patient self-management problems. After the patients demonstrate capability in a behavior, the CDEs proceed to the next behavior. The CDEs continue this process until all potential changes are made to achieve better metabolic control. When the patients actively provide information and ask questions about self-management, the CDEs are likely to provide additional education to address the patients' needs. Typically, the CDEs provide education on and evaluate 1 to 2 behaviors per visit. The number and the distribution of the behavioral assessments are an indicator of underlying patient problems. It is, therefore, meaningful to explore the patterns of and changes in patient problems to provide more adequate health care support.

Diabetes Care at National Taiwan University Hospital

In 2001, the shared care network was implemented at the National Taiwan University Hospital (NTUH), a 2300-bed educational medical center in Taiwan. Currently, the DSME and assessments are documented in the disease management information system (DMIS), developed in 2011 [27]. The DMIS is based on the AADE7 framework and includes the following items: healthy eating, being active, monitoring, taking medication, problem solving, healthy coping, and reducing risks. The detailed contents of the AADE7 are shown in [Multimedia Appendix 1](#). In addition to the shared care network, a telehealthcare program was initiated in 2011 to explore the effectiveness of information technology interventions for individuals with diabetes.

Telehealthcare allows for the promotion of health care services outside the medical institute. Through the combination of information technology and commercial biosensor devices, telehealthcare facilitates longitudinal health status monitoring at a distance [5,14,28,29]. The effectiveness of telehealthcare has been indicated by the industry and policymakers [29-34], and online informatics applications have been widely adopted in diabetes care [1,6,35-37]. It is a promising technology for self-management and can increase patient knowledge, patient

engagement, condition monitoring, long-term follow-ups [1,7,8,16,38], and improved patient outcomes [15,16,38,39].

Methods

Overview

In addition to the routine care of individuals with diabetes mellitus, this research provided a telehealthcare program and aimed to illustrate the way patients used and interacted with the online diabetes self-management system, the impact of a telehealthcare program among 7 self-care skills, and changes in patients' HbA_{1c} levels. A diabetes telehealthcare program was conducted for 18 months, wherein patients received assistance from an online diabetes self-management system to record and manage their daily activities, a 3G glucometer to monitor their glucose, and a teleconsultant service to enhance patients' self-management activities. Behavioral assessments and HbA_{1c} levels were documented in the DMIS. We compared these parameters for those who were assisted by the telehealthcare service (intervention group) at different time points. We also compared the intervention group results to those of patients who did not participate in the telehealthcare program (control group).

The Online Diabetes Self-Management System and Teleconsultant Service


The design of the online diabetes self-management system was based on personal health record (PHR) criteria. The PHR is a health record in which patients' health data and personal information are recorded and maintained by the patients themselves [40]. It is owned, controlled, and managed by the patients [41-43], can be easily accessed, is not limited to

operating systems or devices [41-44], ensures interoperability of data between diverse systems [41,42,44-47], and facilitates easy ways of uploading information [42,43,48]. The PHR emphasizes patient privacy, providing adequate data encryption and a secured environment [38,41-44]. In addition, it improves communication and information sharing between patients and care providers [1,6,41-44,48].

The online diabetes self-management system was developed using C# programming language and a Microsoft SQL server. It was integrated with an off-the-shelf 3G glucometer [14]. The infrastructure design of the online diabetes self-management system is shown in Figure 1. According to the scheme mentioned above, each patient is assigned a unique identification and password, which allows them to log in, edit, and manage their information. The online diabetes self-management system is a Web-based system accessible on the Internet and not limited to operating systems or devices. It adopted the continuity of care document (CCD) standard to enhance the interoperability of data. The 3G glucometer facilitated easy and automatic data uploading after each measurement anywhere and combined blood pressure, blood glucose, and heart rate measurements in 1 instrument.

Encryptions were used to ensure patient privacy and for data transmission. The Web connection between a user client and the server was encrypted using the hypertext transfer protocol secure (HTTPS) protocol, and data transmission between the glucometers and the server was enciphered with advanced encryption standard (AES) encryption. Asynchronous JavaScript and XML (Ajax) and JQuery JavaScript were used to validate the format of the data input and to cooperate with different browsers.

Figure 1. Infrastructure design of the online diabetes self-management system. Patients record their daily activities through manual input, uploading photographs, and a 3G glucometer. The data are transmitted to the system through hypertext transfer protocol secure (HTTPS) and advanced encryption standard (AES) encryption. Caregivers can also enter the system and provide information and support. The system includes information to support self-management. The data can be generated in continuity of care document (CCD) format to ensure the interoperability between systems.



Asynchronous text messages were provided in the online diabetes self-management system; patients and caregivers could communicate through the online diabetes self-management system internal message service or short message service (SMS) text messaging. Application programming interfaces (API) from the Internet were used to send and retrieve information such as weather and pollution standard indexes. The online diabetes self-management system exchanged data with the hospital information system (HIS) using a service-oriented architecture (SOA) mechanism. The Health Level 7 (HL7) embedded Extensible Markup Language (XML) formatted data were used in the framework for data exchange. Patients were able to see their blood test results from the online diabetes self-management system.

The online diabetes self-management system included the monitoring items and the diabetes-related information, such as blood glucose, blood pressure, heart rate, body weight, insulin injection, daily diet, and daily physical activities. Information that was measured with equipment that did not have transmission networks required manual input. Dietary intake could be recorded through the use of either text or images. Additional information to enable self-management and goal setting for glucose control were generated (eg, the mean, median, standard deviation, and maximum and minimum daily blood glucose values). The variations in blood glucose and other parameters are presented together graphically to enable the user to observe the effect of each behavior. The frequency of self-monitoring of blood glucose (SMBG) was recorded and compared with the set goals to determine whether adjustments were needed. Body mass index (BMI) was calculated, and the suggested calorie intake and ingredient volume for each meal were displayed. An additional care-provider interface was designed so that caregivers could get a quick overview of patient status. Case managers were able to log in and view the data uploaded by the patients, identify abnormal events, and make phone calls. The online diabetes self-management system sent an SMS text message to care providers when the data exceeded the alerting range.

This study provided a teleconsultant service to support patients with diabetes self-management. The case managers for this study, including a nurse and a dietitian, were the care providers who interacted with the patients from a distance. They were responsible for monitoring patient status, answering questions about self-care activities, regularly keeping in touch with the patients through telephone calls or text messages, and encouraging them to perform self-management. The care plans and goal setting were formulated through a discussion with each patient during his or her enrollment. The case managers monitored the data uploaded by the patients. They gave advice and reminded the patients to perform self-care activities. In this study, the case managers were not involved in medication adjustments. They did, however, collate patient data and bring the information to the clinic when the patient returned for an appointment. They communicated with physicians to suggest adjustments when needed.

The assistance of the online diabetes self-management system and the teleconsultant service covered the 7 behaviors of self-management activities (shown in [Multimedia Appendix 1](#)).

During the study period, the CDEs did not know whether the patient attended the telehealthcare program.

Patient Enrollment and Data Analysis

Candidates for the study included patients diagnosed with either Type I diabetes mellitus (T1DM) or Type II diabetes mellitus (T2DM) and those with an HbA_{1c} level greater than 7.5 or identified as not well controlled. Patients with severe diabetes complications, such as diabetic foot, diabetic proliferative retinopathy, liver dysfunction, end-stage renal disease, or other medical problems that could affect the study results or trial participation were excluded. The patients were approached during regular follow-up visits in their physician's office, and informed consent was obtained from each participant. After enrollment, patients were taught to use the online diabetes self-management system and the glucometer, and they were informed how to contact the case manager for assistance. The glucometer and test strips were provided for glucose monitoring without a charge. Patients were allowed to use their own glucometer if they preferred. Those who chose not to use the provided glucometer were allowed to input their data manually. This study was reviewed and approved by the NTUH Institutional Review Board (IRB) (No. 201108018RC).

By the end of February 2013, 184 patients were enrolled in the telehealthcare program. Of the patients who participated in the telehealthcare program, 59 (32.1%) were also in the shared care network. The results of participants in the telehealthcare and shared care network were also compared to those of the participants who did not use the telehealthcare service but did complete the behavioral assessments and laboratory monitoring; these individuals served as the control participants. We recruited 103 control participants and matched their demographic characteristics with those of the telehealthcare participants to minimize the effect of potential confounding variables, including gender, age, diabetes type, duration, years of participation in the shared care network, and insulin treatment. We focused on analyzing the data of patients in the telehealthcare and the control groups, who were evaluated by behavioral assessments and laboratory monitoring every 3 months.

The evaluation of this study consisted of 3 sections (shown in [Figure 2](#)). We first analyzed the logged records of the system and illustrated the way patients used the online diabetes self-management system. For the second and third sections, the behavioral assessments and laboratory results from the DMIS for 18 months were evaluated by within-group comparisons at different time points for those participants who entered the telehealthcare program. We also compared the telehealthcare and control groups at different time points.

The data were grouped into several time ranges (see [Figure 3](#)). The time ranges included the entire 18 months of patient participation (September 2011 to February 2013, T1), the first 6 months (September 2011 to February 2012, T2), the second 6 months (March 2012 to August 2012, T3), and the last 6 months of patient participation (September 2012 to February 2013, T4). The data were grouped in 6-month intervals to obtain 1 or 2 records from regular patient visits.

The second section was the behavioral analysis. The number and distribution of the behavioral assessments indicated the problems of patients while performing self-management. In the behavioral analysis, the number of behavioral assessments was calculated for the 7 behaviors for each time range. The number of behavioral assessments of the telehealthcare participants during T1 was compared with each range (T2, T3, and T4) to demonstrate the overall variation of each time range, and the baseline (T2) was compared with T3 and T4 to see the variation across time. The differences between the telehealthcare and the control participants at each range (T1, T2, T3, and T4) were also compared.

The third section was the HbA_{1c} analysis, which consisted of calculating the HbA_{1c} mean and analyzing HbA_{1c} variability. HbA_{1c} variability represents intraindividual differences for each patient, which refers to the changes in glycemia over longer

periods of time reflected in changes in HbA_{1c} from one visit to the next [49,50]. It is defined as the standard deviation (SD) of the serial HbA_{1c} measurements. All data for blood tests within each time range were collected, and the HbA_{1c} mean and HbA_{1c} variability were calculated for each patient during each time range. The value for the telehealthcare participants at each time range (T2, T3, and T4) was compared to T1 to observe the overall variation of each time range, and the changes at T3 and T4 were observed by comparing to their baselines (T2). The differences between the telehealthcare and control participants at each time range (T1, T2, T3, and T4) were compared.

Paired *t* tests were used to analyze the differences in variables of interest between the time points of the telehealthcare participants' measures. Independent *t* tests were used to compare the telehealthcare and control results and between T1 and each time range for telehealthcare participants. We used SPSS version 17.0 (SPSS Inc, Chicago, IL, USA) for the statistical analyses.

Figure 2. A flowchart of the data analysis.





Figure 3. Analysis timeline.



Results

Table 1 and **Multimedia Appendices 2-4** show the demographic information of the telehealthcare and control participants, including gender, age, years of participation in the shared care network, disease duration, and the use of insulin injection. The telehealthcare group consisted of 18 T1DM and 41 T2DM patients. The control group consisted of 32 T1DM and 71 T2DM patients. The average age of the telehealthcare group was 51.34 years (SD 12.79). **Figure 4** shows the user interface of the online diabetes self-management system. Overall, 90% (53/59) of the patients logged in and used the online diabetes self-management system. On average, patients logged in 1.3 (SD 2.2) times every week and performed 1.1 (SD 1.3) SMBG daily. Analysis of the self-management records showed that 98% (58/59) of the patients documented blood glucose, 73% (43/59) documented blood pressure, 69% (41/59) documented heart rate, 44% (26/59) documented dietary record, 44% (26/59) documented insulin injections, and 31% (31/59) documented physical activities. In all, 61% (36/59) of the patients used telephone services to contact the case managers; 56% (33/59) used text messages.

Table 2 shows the behavioral assessments of telehealthcare group during different time ranges. SMBG was significantly

different between T1 and T4 ($P<.001$) and T2 and T4 ($P=.046$). Most of the number of assessments increased at T3 and decreased at T4, and SMBG was the only behavior that continued to increase until T4. There was a statistically significant difference between the telehealthcare and control participants in 5 behaviors at T1 (**Table 3**), including being active ($P<.001$), healthy eating ($P<.001$), taking medication ($P<.001$), healthy coping ($P=.02$), and problem solving ($P<.001$). SMBG was significantly different between the telehealthcare and control groups at T3 ($P=.02$). Notably, the telehealthcare participants had more assessments in healthy coping and reducing risk at T4 than did the control participants.

The HbA_{1c} mean and variability differences for the telehealthcare group are shown in **Table 4**. The mean HbA_{1c} level decreased significantly ($P=.02$) at T3 compared to the baseline (T2) and slightly increased at T4, and the HbA_{1c} mean decreased from 7.8 to 7.68. There was a significant difference in HbA_{1c} variability between T1 and T2 ($P<.001$), T1 and T3 ($P<.001$), and T1 and T4 ($P<.001$), and the HbA_{1c} variability decreased from 0.30 to 0.23 during patient participation. The HbA_{1c} mean and variability differences between the telehealthcare and control groups are shown in **Table 5**, and were not statistically significant.

Table 1. Patient demographics, including gender, age, and diabetes type (N=162).

| Demographic information | Telehealthcare n=59 | Control n=103 |
|-------------------------|------------------------|------------------|
| Gender, n (%) | | |
| Male | 29 (49.2) | 52 (50.5) |
| Female | 30 (50.8) | 51 (49.5) |
| Age, mean (SD) | 51.3 (12.8) | 52.55 (12.19) |
| T1DM^a | | |
| <45 years, n (%) | 10 (55.6) | 23 (71.9) |
| >45 years, n (%) | 8 (44.4) | 9 (28.1) |
| Age, mean (SD) | 41.78 (9.78) | 41.25 (8.16) |
| T2DM^b | | |
| <65 years, n (%) | 32 (78.0) | 53 (74.6) |
| >65 years, n (%) | 9 (22.0) | 18 (25.4) |
| Age, mean (SD) | 55.54 (11.73) | 57.65 (10.11) |

^aT1DM: Type 1 diabetes mellitus.

^bT2DM: Type 2 diabetes mellitus.

Table 2. The American Association of Diabetes Educators 7 Self-Care Behaviors (AADE7) education of patients with the telehealthcare service (n=59).

| Time ^a and AADE7 behavior | Behavioral assessments, mean (SD) | Comparison with T1, <i>P</i> value | Comparison with T2, <i>P</i> value |
|--------------------------------------|-----------------------------------|------------------------------------|------------------------------------|
| T1 (n=59) | | | |
| AADE7 education | | | |
| Being active | 0.36 (0.36) | | |
| Healthy eating | 0.60 (0.35) | | |
| Taking medication | 0.51 (0.36) | | |
| Healthy coping | 0.09 (0.22) | | |
| Problem solving | 0.31 (0.36) | | |
| Reducing risks | 0.02 (0.10) | | |
| Monitoring | 0.56 (0.15) | | |
| T2 (n=59) | | | |
| Being active | 0.34 (0.54) | .66 | |
| Healthy eating | 0.58 (0.59) | .66 | |
| Taking medication | 0.53 (0.60) | .95 | |
| Healthy coping | 0.08 (0.28) | .90 | |
| Problem solving | 0.42 (0.53) | .18 | |
| Reducing risks | 0.00 (0.00) | .10 | |
| Monitoring | 0.63 (0.69) | .68 | |
| T3 (n=59) | | | |
| Being active | 0.42 (0.56) | .44 | .42 |
| Healthy eating | 0.66 (0.63) | .55 | .49 |
| Taking medication | 0.54 (0.60) | .71 | .89 |
| Healthy coping | 0.10 (0.30) | .82 | .66 |
| Problem solving | 0.27 (0.49) | .62 | .07 |
| Reducing risks | 0.03 (0.26) | .76 | .32 |
| Monitoring | 0.68 (0.71) | .23 | .73 |
| T4 (n=59) | | | |
| Being active | 0.31 (0.60) | .58 | .86 |
| Healthy eating | 0.58 (0.60) | .75 | .27 |
| Taking medication | 0.46 (0.63) | .59 | .90 |
| Healthy coping | 0.08 (0.28) | .90 | .90 |
| Problem solving | 0.24 (0.54) | .38 | .18 |
| Reducing risks | 0.03 (0.18) | .68 | .16 |
| Monitoring | 0.75 (0.63) | <.001 | .046 |

^aT1: Entire duration of the telehealthcare program, from September 2011 to February 2013 (18 months); T2: initial stage of the telehealthcare program, from September 2011 to February 2012 (6 months); T3: middle stage of the telehealthcare program, from March 2012 to August 2012 (6 months); T4: last stage of the telehealthcare program, from September 2012 to February 2013 (6 months); T4: Last stage of the telehealthcare program, from September 2012 to February 2013 (6 months).

Table 3. The American Association of Diabetes Educators 7 Self-Care Behaviors (AADE7) education of patients with and without the telehealthcare service.

| Time ^a and AADE7 behavior | Behavioral assessments per patient, mean (SD) | | P value |
|--------------------------------------|---|--------------------|---------|
| | Telehealthcare (n=59) | Control (n=103) | |
| T1 | | | |
| Being active | 0.36 (0.36) | 0.44 (0.41) | <.001 |
| Healthy eating | 0.60 (0.35) | 0.72 (0.43) | <.001 |
| Taking medication | 0.51 (0.36) | 0.50 (0.40) | <.001 |
| Healthy coping | 0.09 (0.22) | 0.08 (0.17) | .02 |
| Problem solving | 0.31 (0.36) | 0.34 (0.37) | <.001 |
| Reducing risks | 0.02 (0.10) | 0.02 (0.11) | .29 |
| Monitoring | 0.56 (0.15) | 0.60 (0.41) | .94 |
| T2 | | | |
| Being active | 0.34 (0.54) | 0.45 (0.61) | .27 |
| Healthy eating | 0.58 (0.59) | 0.71 (0.67) | .20 |
| Taking medication | 0.53 (0.60) | 0.49 (0.63) | .69 |
| Healthy coping | 0.08 (0.28) | 0.08 (0.31) | .91 |
| Problem solving | 0.42 (0.53) | 0.45 (0.70) | .84 |
| Reducing risks | 0.00 (0.00) | 0.01 (0.10) | .45 |
| Monitoring | 0.63 (0.69) | 0.80 (0.74) | .14 |
| T3 | | | |
| Being active | 0.42 (0.56) | 0.42 (0.57) | .93 |
| Healthy eating | 0.66 (0.63) | 0.71 (0.73) | .65 |
| Taking medication | 0.54 (0.60) | 0.52 (0.67) | .87 |
| Healthy coping | 0.10 (0.30) | 0.10 (0.30) | .96 |
| Problem solving | 0.27 (0.49) | 0.31 (0.50) | .66 |
| Reducing risks | 0.03 (0.26) | 0.02 (0.14) | .66 |
| Monitoring | 0.68 (0.71) | 0.42 (0.53) | .02 |
| T4 | | | |
| Being active | 0.31 (0.60) | 0.48 (0.64) | .11 |
| Healthy eating | 0.58 (0.59) | 0.73 (0.70) | .18 |
| Taking medication | 0.46 (0.63) | 0.50 (0.61) | .64 |
| Healthy coping | 0.08 (0.28) | 0.05 (0.22) | .38 |
| Problem solving | 0.24 (0.54) | 0.25 (0.44) | .80 |
| Reducing risks | 0.03 (0.18) | 0.02 (0.20) | .66 |
| Monitoring | 0.75 (0.63) | 1.69 (0.47) | .13 |

^aT1: The entire duration of the telehealthcare program, from September 2011 to February 2013 (18 months); T2: Initial stage of the telehealthcare program, from September 2011 to February 2012 (6 months); T3: Middle stage of the telehealthcare program, from March 2012 to August 2012 (6 months); T4: Last stage of the telehealthcare program, from September 2012 to February 2013 (6 months).

Table 4. The HbA_{1c} mean and variability of the patients with the telehealthcare service (n=59).

| Time ^a and HbA _{1c} | Mean (SD) | Compared with T1, <i>P</i> value | Compared with T2, <i>P</i> value |
|---|-------------|----------------------------------|----------------------------------|
| T1 | | | |
| Mean | 7.72 (0.51) | | |
| Variability | 0.51 (0.29) | | |
| T2 | | | |
| Mean | 7.80 (0.38) | .48 | |
| Variability | 0.30 (0.31) | <.001 | |
| T3 | | | |
| Mean | 7.64 (0.40) | .62 | .02 |
| Variability | 0.31 (0.31) | <.001 | .81 |
| T4 | | | |
| Mean | 7.68 (0.31) | .80 | .17 |
| Variability | 0.23 (0.18) | <.001 | .11 |

^aT1: The entire duration of the telehealthcare program, from September 2011 to February 2013 (18 months); T2: Initial stage of the telehealthcare program, from September 2011 to February 2012 (6 months); T3: Middle stage of the telehealthcare program, from March 2012 to August 2012 (6 months);

Table 5. The HbA_{1c} mean and variability of the patients with and without the telehealthcare service.

| Time ^a and HbA _{1c} | HbA _{1c} , mean (SD) | | <i>P</i> value |
|---|-------------------------------|--------------------|----------------|
| | Telehealthcare (n=59) | Control (n=103) | |
| T1 | | | |
| Mean | 7.72 (0.51) | 7.65 (0.52) | .70 |
| Variability | 0.51 (0.29) | 0.52 (0.32) | .78 |
| T2 | | | |
| Mean | 7.80 (0.38) | 7.67 (1.45) | .39 |
| Variability | 0.30 (0.31) | 0.23 (0.27) | .14 |
| T3 | | | |
| Mean | 7.64 (0.40) | 7.67 (1.57) | .89 |
| Variability | 0.31 (0.31) | 0.27 (0.26) | .45 |
| T4 | | | |
| Mean | 7.68 (0.31) | 7.66 (0.34) | .94 |
| Variability | 0.23 (0.18) | 0.25 (0.23) | .46 |

^aT1: The entire duration of the telehealthcare program, from September 2011 to February 2013 (18 months); T2: Initial stage of the telehealthcare program, from September 2011 to February 2012 (6 months); T3: Middle stage of the telehealthcare program, from March 2012 to August 2012 (6 months); T4: Last stage of the telehealthcare program, from September 2012 to February 2013 (6 months).

Figure 4. Screenshot of the online diabetes self-management system user interface (note that the original user interface is in Chinese).

| New Edit Delete | | |
|---------------------|---------------|--------------|
| Time of Measurement | Blood Glucose | Glucose Type |
| 2012/12/05 14:37:00 | 128 | |
| 2012/12/05 12:38:00 | 93 | AC |
| 2012/12/04 22:39:00 | 104 | PC |
| 2012/12/04 06:30:00 | 96 | AC |
| 2012/12/03 22:43:00 | 168 | PC |
| 2012/12/03 20:43:00 | 131 | AC |
| 2012/12/03 08:27:00 | 115 | AC |
| 2012/12/02 20:39:00 | 127 | PC |
| 2012/12/02 10:19:00 | 126 | AC |
| 2012/12/01 20:55:00 | 111 | PC |

Discussion

Principal Findings

The aims of this study were to determine how diabetic patients use online applications, determine the impact of the telehealthcare program on the patients' 7 self-care behaviors, and examine HbA_{1c} level changes during the course of participation in such a program. Through the use of the online diabetes self-management system and the teleconsultant service, the diabetic patients managed complex information about their diseases and obtained support from their care providers. Patients were reminded to perform SMBG, and their self-care behaviors were reinforced. In all, 90% of the patients used the online diabetes self-management system, and 98% performed SMBG. On average, patients logged in every week and performed the SMBG daily. Patients who maintain insulin records are also likely to maintain dietary records, which are related to the effects of insulin injections on food intake. In this study, the use of an image uploading application did not cause more patients to maintain dietary records. One explanation for this finding is that there were more elderly patients in this program and these patients do not often use smartphones or carry a camera with them.

The case managers observed that technological difficulties were the main reason for a decline in the use of the online diabetes self-management system by elderly patients. Some of the patients claimed that they were too busy to use the online diabetes self-management system. Previous research pointed out that elderly people had poor technical skills [41]. In our study, the average age of the enrolled patients was older than 50 years; blood glucose, blood pressure, and heart rate were the 3 items most often recorded. This indicates that a proportion of

the patients only used the 3G glucometer to upload data, but seldom logged into the online diabetes self-management system. Despite the technological difficulties, patients unfamiliar with information technologies were not troubled with the computer applications used in this program, and they used the 3G glucometer to participate and obtain support from their case managers in a seamless manner. Easier ways to upload data may increase the participation of those unfamiliar with technology, enhance the completeness of the dataset, and allow them to obtain support from a distance.

According to the online diabetes self-management system logged records, more patients used the phone call service than the messaging service to contact the case managers. However, the case managers found that the younger patients used text messages more often than the elderly patients did. They used various applications, such as email, smartphone applications (eg, Whatsapp, LINE), and SMS text messages, to communicate with patients in order to accommodate the patients' busy lifestyles, which made the records difficult to trace and unavailable for presentation here. Some of the patients expressed a reluctance to let their colleagues or the people around them know about their disease, and they considered the asynchronous messages useful in allowing them to communicate with caregivers without the risk of being overheard by others. Most of the elderly patients were not familiar with keyboard typing, and they relied on phone calls to communicate. It is worth noting that providing free test strips or services created an incentive for patient participation. In fact, most of the patients were reluctant to continue their participation when told the test strips or the service may no longer be free, indicating the financial burden borne by persons with diabetes.

In the behavioral analysis, the number of SMBG assessments increased significantly in the last stage, and the patients that participated in the telehealthcare service had significantly more SMBG assessments than did those without the telehealthcare service in the 6-12 month stage. Because the telehealthcare service used was based on glucose measurements, the case managers provided support and suggestions after they observed the uploaded data. SMBG was the primary self-care ability enhanced through participation in the telehealthcare program. By performing SMBG more regularly, patients encountered problems and discussed these problems more often with the CDEs.

When the patients who participated in the telehealthcare program for 18 months were compared to those who did not, 5 of the 7 behaviors showed significant differences. Although the CDEs enhance patient education based on different considerations, patients in the telehealthcare program required less support in being active, healthy eating, and problem solving, and required more support in taking medicine and healthy coping. One explanation is that the skill of insulin injection and the overcoming of psychological obstacles still requires face-to-face interaction. Each visit was done with 1 or 2 behavior assessments, and did not cover all 7 behaviors. Some of the behaviors had not been assessed yet, and were unlikely to show differences during the 6-month period. The overall time frame concluded the 7 behaviors and also represented more time to see the changes after the education. During the last period of patient participation, the assessments of healthy coping and reducing risks were higher for those who participated in the telehealthcare program compared to those who did not. Healthy coping and reducing risks are the skills that the CDEs enhanced when the patients demonstrated that they were capable of coping with their other problems. The observation of an increase in these 2 behaviors implies that the patients were more skillful.

The mean value of HbA_{1c} of participants at 6 to 12 months improved significantly compared to baseline, and slightly declined in the last period. This result indicates that the patients experienced a “worn out” period. When the patients first entered the program, they were more conscious of their self-management behaviors because they knew that someone was watching them; as a result, they improved significantly. However, after the patients became more familiar with the service and were less anxious about the program, a slight decline was observed. The HbA_{1c} variability of each 6-month interval was significantly lower than the overall HbA_{1c} variability. The decrease of the mean HbA_{1c} value implies that patients who participated in the program improved substantially across the 18-month period, and have potentially reduced the risks of complication development with less HbA_{1c} variability [49,50]. The HbA_{1c} mean demonstrates the overall glycemic control in a 3-month period, and is limited to show the variation of glycemic control during each time interval. The behaviors affected the performance of HbA_{1c}, and the change of behaviors requires time to show its effect. Therefore, there may not be differences during each 6-month range, but significant differences in the long term.

Limitations

In this small-scale pilot study, we provided a telehealthcare program that consisted of 3G glucometers, free test strips, an online diabetes self-management system, and easy access to professional support. All the program components contributed to patient improvements, although we did not measure the individual contribution of each of the components. Unfortunately, the contribution of each of the components remains unclear; this is a limitation of this study. However, those with T1DM received free test strips from the NHI. The program did produce a small benefit of HbA_{1c} control for the T1DM patients (HbA_{1c} decreased from 7.83 to 7.74), implying that the free glucometer strips were not solely responsible for the outcomes of this program.

The mean HbA_{1c} value for the telehealthcare group during the last period dropped to 7.68 and was still considered not well controlled. This may be because of the enrollment of patients with very poor glycemic control. In addition, the therapeutic responses may require more time since HbA_{1c} is a measure of average blood glucose over the course of 3 months. The patient education in this study was based on the observations of different CDEs (3 nurses and 2 dietitians), and the evaluation result may differ from CDE to CDE. Another limitation was that before the development of the DMIS, the documentation of patient education was paper-based rather than structured in the AADE7 form. The DMIS went online in July 2011 and stabilized in August 2011; the telehealthcare program was initiated in September 2011. Hence, we were unable to obtain patients' documentation before their entry into the telehealthcare program.

While connecting the SMBG assessment and the performance of HbA_{1c} level, it could be observed that the number of SMBG assessments increased significantly and the mean HbA_{1c} level slightly increased in the last stage of patient participation. The number of SMBG assessments may refer to patients performing more SMBG and may also refer to CDEs trying to identify patient problems through encouraging them to perform more SMBG. This study has not further explored the reason of the increasing of SMBG assessment; hence, it could not explain the reason for the increasing of SMBG assessments. Further research is needed to measure the contribution of each component of the telehealthcare program and determine how to improve patient performance when they are worn out. More work is needed to demonstrate the effect of telehealthcare on specific behaviors.

Conclusions

This study showed that using a sophisticated technological design supported the patients with diabetes in self-management. It appears that telehealthcare is effective in enhancing blood glucose monitoring, and the patients in the program showed improvements in glycemic control. The self-care behaviors affected patient outcomes and the changes in behavior required time to show effects. Telehealthcare has a positive effect on patients with diabetes, and it may encourage more technological interventions for diabetes care.

Acknowledgments

We thank the National Science Council of Taiwan for funding the program (NSC 101-2219-E-002-024). We are thankful for the NTUH IRB approval (No. 201108018RC). We express our gratitude to the cooperation of the Diabetes Center and Information Systems Office of NTUH, and the participation of the educators and the case managers. The initial concept of the system architecture of the online diabetes self-management system has been accepted by the Asian-Pacific Chinese Diabetes Forum (2013) in Chinese.

Conflicts of Interest

None declared.

Multimedia Appendix 1

The detailed contents of the American Association of Diabetes Educators 7 Self-Care Behaviors (AADE7) education and the telehealthcare service.

[\[PDF File \(Adobe PDF File\), 29KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Patient demographics, years of participation in the shared care network.

[\[PDF File \(Adobe PDF File\), 23KB-Multimedia Appendix 2\]](#)

Multimedia Appendix 3

Patient demographics, duration of diabetes.

[\[PDF File \(Adobe PDF File\), 22KB-Multimedia Appendix 3\]](#)

Multimedia Appendix 4

Patient demographics, insulin injection frequency.

[\[PDF File \(Adobe PDF File\), 23KB-Multimedia Appendix 4\]](#)

References

1. Lin YH, Chen RR, Guo SH, Chang HY, Chang HK. Developing a web 2.0 diabetes care support system with evaluation from care provider perspectives. *J Med Syst* 2012 Aug;36(4):2085-2095. [doi: [10.1007/s10916-011-9672-7](https://doi.org/10.1007/s10916-011-9672-7)] [Medline: [21369781](https://pubmed.ncbi.nlm.nih.gov/21369781/)]
2. Glasgow RE, Christiansen SM, Kurz D, King DK, Woolley T, Faber AJ, et al. Engagement in a diabetes self-management website: usage patterns and generalizability of program use. *Journal of Medical Internet Research* 2011;13(1):e9. [doi: [10.2196/jmir.1391](https://doi.org/10.2196/jmir.1391)]
3. Bourbeau J, Nault D, Dang-Tan T. Self-management and behaviour modification in COPD. *Patient Educ Couns* 2004 Mar;52(3):271-277. [doi: [10.1016/S0738-3991\(03\)00102-2](https://doi.org/10.1016/S0738-3991(03)00102-2)] [Medline: [14998597](https://pubmed.ncbi.nlm.nih.gov/14998597/)]
4. Bodenheimer T, Lorig K, Holman H, Grumbach K. Patient self-management of chronic disease in primary care. *JAMA* 2002 Nov 20;288(19):2469-2475. [Medline: [12435261](https://pubmed.ncbi.nlm.nih.gov/12435261/)]
5. Heinemann L. Measuring glucose concentrations: daily practice, current and future developments. *J Diabetes Sci Technol* 2008 Jul;2(4):710-717 [FREE Full text] [Medline: [19885250](https://pubmed.ncbi.nlm.nih.gov/19885250/)]
6. Watson AJ, Kvedar JC, Rahman B, Pelletier AC, Salber G, Grant RW. Diabetes connected health: a pilot study of a patient- and provider-shared glucose monitoring web application. *J Diabetes Sci Technol* 2009 Mar;3(2):345-352 [FREE Full text] [Medline: [20144366](https://pubmed.ncbi.nlm.nih.gov/20144366/)]
7. Klonoff DC. Using telemedicine to improve outcomes in diabetes--an emerging technology. *J Diabetes Sci Technol* 2009 Jul;3(4):624-628 [FREE Full text] [Medline: [20144303](https://pubmed.ncbi.nlm.nih.gov/20144303/)]
8. Grant RW, Wald JS, Poon EG, Schnipper JL, Gandhi TK, Volk LA, et al. Design and implementation of a web-based patient portal linked to an ambulatory care electronic health record: patient gateway for diabetes collaborative care. *Diabetes Technol Ther* 2006 Oct;8(5):576-586. [doi: [10.1089/dia.2006.8.576](https://doi.org/10.1089/dia.2006.8.576)] [Medline: [17037972](https://pubmed.ncbi.nlm.nih.gov/17037972/)]
9. Whittemore R, Jaser SS, Faulkner MS, Murphy K, Delamater A, Grey M, TEENCOPE Research Group. Type 1 diabetes eHealth psychoeducation: youth recruitment, participation, and satisfaction. *J Med Internet Res* 2013;15(1):e15 [FREE Full text] [doi: [10.2196/jmir.2170](https://doi.org/10.2196/jmir.2170)] [Medline: [23360729](https://pubmed.ncbi.nlm.nih.gov/23360729/)]
10. Wang J, Zgibor J, Matthews JT, Charron-Prochownik D, Sereika SM, Siminerio L. Self-monitoring of blood glucose is associated with problem-solving skills in hyperglycemia and hypoglycemia. *Diabetes Educ* 2012;38(2):207-218. [doi: [10.1177/0145721712440331](https://doi.org/10.1177/0145721712440331)] [Medline: [22454405](https://pubmed.ncbi.nlm.nih.gov/22454405/)]

11. Zgibor JC, Peyrot M, Ruppert K, Noulet W, Siminerio LM, Peeples M, AADE/UPMC Diabetes Education Outcomes Project. Using the American Association of Diabetes Educators Outcomes System to identify patient behavior change goals and diabetes educator responses. *Diabetes Educ* 2007;33(5):839-842. [doi: [10.1177/0145721707307611](https://doi.org/10.1177/0145721707307611)] [Medline: [17925588](https://pubmed.ncbi.nlm.nih.gov/17925588/)]
12. Funnell MM, Brown TL, Childs BP, Haas LB, Hoseney GM, Jensen B, et al. National standards for diabetes self-management education. *Diabetes Care* 2012 Jan;35 Suppl 1:S101-S108 [FREE Full text] [doi: [10.2337/dc12-s101](https://doi.org/10.2337/dc12-s101)] [Medline: [22187467](https://pubmed.ncbi.nlm.nih.gov/22187467/)]
13. Norris SL, Lau J, Smith SJ, Schmid CH, Engelgau MM. Self-management education for adults with type 2 diabetes: a meta-analysis of the effect on glycemic control. *Diabetes Care* 2002 Jul;25(7):1159-1171. [Medline: [12087014](https://pubmed.ncbi.nlm.nih.gov/12087014/)]
14. Chen LC, Chen CW, Weng YC, Shang RJ, Yu HC, Chung Y, et al. An information technology framework for strengthening telehealthcare service delivery. *Telemed J E Health* 2012 Oct;18(8):596-603 [FREE Full text] [doi: [10.1089/tmj.2011.0267](https://doi.org/10.1089/tmj.2011.0267)] [Medline: [23061641](https://pubmed.ncbi.nlm.nih.gov/23061641/)]
15. Lim S, Kang SM, Shin H, Lee HJ, Won Yoon J, Yu SH, et al. Improved glycemic control without hypoglycemia in elderly diabetic patients using the ubiquitous healthcare service, a new medical information system. *Diabetes Care* 2011 Feb;34(2):308-313 [FREE Full text] [doi: [10.2337/dc10-1447](https://doi.org/10.2337/dc10-1447)] [Medline: [21270188](https://pubmed.ncbi.nlm.nih.gov/21270188/)]
16. Hess R, Bryce CL, Paone S, Fischer G, McTigue KM, Olshansky E, et al. Exploring challenges and potentials of personal health records in diabetes self-management: implementation and initial assessment. *Telemed J E Health* 2007 Oct;13(5):509-517. [doi: [10.1089/tmj.2006.0089](https://doi.org/10.1089/tmj.2006.0089)] [Medline: [17999613](https://pubmed.ncbi.nlm.nih.gov/17999613/)]
17. Hieftje K, Edelman EJ, Camenga DR, Fiellin LE. Electronic media-based health interventions promoting behavior change in youth: a systematic review. *JAMA Pediatr* 2013 Jun;167(6):574-580 [FREE Full text] [doi: [10.1001/jamapediatrics.2013.1095](https://doi.org/10.1001/jamapediatrics.2013.1095)] [Medline: [23568703](https://pubmed.ncbi.nlm.nih.gov/23568703/)]
18. Pal K, Eastwood SV, Michie S, Farmer AJ, Barnard ML, Peacock R, et al. Computer-based diabetes self-management interventions for adults with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2013;3:CD008776. [doi: [10.1002/14651858.CD008776.pub2](https://doi.org/10.1002/14651858.CD008776.pub2)] [Medline: [23543567](https://pubmed.ncbi.nlm.nih.gov/23543567/)]
19. Lin RL, Lin FJ, Wu CL, Peng MJ, Chen PJ, Kuo HT. Effect of a hospital-based case management approach on treatment outcome of patients with tuberculosis. *J Formos Med Assoc* 2006 Aug;105(8):636-644. [doi: [10.1016/S0929-6646\(09\)60162-5](https://doi.org/10.1016/S0929-6646(09)60162-5)] [Medline: [16935764](https://pubmed.ncbi.nlm.nih.gov/16935764/)]
20. Ofman JJ, Badamgarav E, Henning JM, Knight K, Gano AD, Levan RK, et al. Does disease management improve clinical and economic outcomes in patients with chronic diseases? A systematic review. *Am J Med* 2004 Aug 1;117(3):182-192. [Medline: [15300966](https://pubmed.ncbi.nlm.nih.gov/15300966/)]
21. Chan DC, Heidenreich PA, Weinstein MC, Fonarow GC. Heart failure disease management programs: a cost-effectiveness analysis. *Am Heart J* 2008 Feb;155(2):332-338. [doi: [10.1016/j.ahj.2007.10.001](https://doi.org/10.1016/j.ahj.2007.10.001)] [Medline: [18215605](https://pubmed.ncbi.nlm.nih.gov/18215605/)]
22. Markle A. The economic impact of case management. *Case Manager* 2004;15(4):54-58. [doi: [10.1016/S1061925904001328](https://doi.org/10.1016/S1061925904001328)] [Medline: [15247897](https://pubmed.ncbi.nlm.nih.gov/15247897/)]
23. Shumway M, Boccellari A, O'Brien K, Okin RL. Cost-effectiveness of clinical case management for ED frequent users: results of a randomized trial. *Am J Emerg Med* 2008 Feb;26(2):155-164. [doi: [10.1016/j.ajem.2007.04.021](https://doi.org/10.1016/j.ajem.2007.04.021)] [Medline: [18272094](https://pubmed.ncbi.nlm.nih.gov/18272094/)]
24. Lai MS. National health insurance and the way leading to better diabetes care in Taiwan. Is there a role of comprehensive analyses of the claims data? *J Formos Med Assoc* 2012 Nov;111(11):587-588. [doi: [10.1016/j.jfma.2012.08.016](https://doi.org/10.1016/j.jfma.2012.08.016)] [Medline: [23217593](https://pubmed.ncbi.nlm.nih.gov/23217593/)]
25. Chang TJ, Jiang YD, Chang CH, Chung CH, Yu NC, Chuang LM. Accountability, utilization and providers for diabetes management in Taiwan, 2000-2009: an analysis of the National Health Insurance database. *J Formos Med Assoc* 2012 Nov;111(11):605-616. [doi: [10.1016/j.jfma.2012.09.011](https://doi.org/10.1016/j.jfma.2012.09.011)] [Medline: [23217596](https://pubmed.ncbi.nlm.nih.gov/23217596/)]
26. Jiang YD, Chang CH, Tai TY, Chen JF, Chuang LM. Incidence and prevalence rates of diabetes mellitus in Taiwan: analysis of the 2000-2009 Nationwide Health Insurance database. *J Formos Med Assoc* 2012 Nov;111(11):599-604. [doi: [10.1016/j.jfma.2012.09.014](https://doi.org/10.1016/j.jfma.2012.09.014)] [Medline: [23217595](https://pubmed.ncbi.nlm.nih.gov/23217595/)]
27. Chen L, Yu HC, Li HC, Wang YV, Chen HJ, Wang IC, et al. An architecture model for multiple disease management information systems. *J Med Syst* 2013 Apr;37(2):9931. [doi: [10.1007/s10916-013-9931-x](https://doi.org/10.1007/s10916-013-9931-x)] [Medline: [23423776](https://pubmed.ncbi.nlm.nih.gov/23423776/)]
28. Wagner EH. The role of patient care teams in chronic disease management. *BMJ* 2000 Feb 26;320(7234):569-572 [FREE Full text] [Medline: [10688568](https://pubmed.ncbi.nlm.nih.gov/10688568/)]
29. McLean S, Protti D, Sheikh A. Telehealthcare for long term conditions. *BMJ* 2011;342:d120. [Medline: [21292710](https://pubmed.ncbi.nlm.nih.gov/21292710/)]
30. Paré G, Jaana M, Sicotte C. Systematic review of home telemonitoring for chronic diseases: the evidence base. *J Am Med Inform Assoc* 2007;14(3):269-277 [FREE Full text] [doi: [10.1197/jamia.M2270](https://doi.org/10.1197/jamia.M2270)] [Medline: [17329725](https://pubmed.ncbi.nlm.nih.gov/17329725/)]
31. Galbreath AD, Krasuski RA, Smith B, Stajduhar KC, Kwan MD, Ellis R, et al. Long-term healthcare and cost outcomes of disease management in a large, randomized, community-based population with heart failure. *Circulation* 2004 Dec 7;110(23):3518-3526 [FREE Full text] [doi: [10.1161/01.CIR.0000148957.62328.89](https://doi.org/10.1161/01.CIR.0000148957.62328.89)] [Medline: [15531765](https://pubmed.ncbi.nlm.nih.gov/15531765/)]
32. Goldzweig CL, Towfigh A, Maglione M, Shekelle PG. Costs and benefits of health information technology: new trends from the literature. *Health Aff (Millwood)* 2009;28(2):w282-w293 [FREE Full text] [doi: [10.1377/hlthaff.28.2.w282](https://doi.org/10.1377/hlthaff.28.2.w282)] [Medline: [19174390](https://pubmed.ncbi.nlm.nih.gov/19174390/)]
33. Costa BM, Fitzgerald KJ, Jones KM, Dunning Am T. Effectiveness of IT-based diabetes management interventions: a review of the literature. *BMC Fam Pract* 2009;10:72 [FREE Full text] [doi: [10.1186/1471-2296-10-72](https://doi.org/10.1186/1471-2296-10-72)] [Medline: [19917136](https://pubmed.ncbi.nlm.nih.gov/19917136/)]

34. Verhoeven F, Tanja-Dijkstra K, Nijland N, Eysenbach G, van Gemert-Pijnen L. Asynchronous and synchronous teleconsultation for diabetes care: a systematic literature review. *J Diabetes Sci Technol* 2010 May;4(3):666-684 [FREE Full text] [Medline: [20513335](#)]
35. Mougiakakou SG, Bartsocas CS, Bozas E, Chaniotakis N, Iliopoulou D, Kouris I, et al. SMARTDIAB: a communication and information technology approach for the intelligent monitoring, management and follow-up of type 1 diabetes patients. *IEEE Trans Inf Technol Biomed* 2010 May;14(3):622-633. [doi: [10.1109/TITB.2009.2039711](#)] [Medline: [20123578](#)]
36. Tani S, Marukami T, Matsuda A, Shindo A, Takemoto K, Inada H. Development of a health management support system for patients with diabetes mellitus at home. *J Med Syst* 2010 Jun;34(3):223-228. [Medline: [20503606](#)]
37. Gómez EJ, Hernando Pérez ME, Vering T, Rigla Cros M, Bott O, García-Sáez G, et al. The INCA system: a further step towards a telemedical artificial pancreas. *IEEE Trans Inf Technol Biomed* 2008 Jul;12(4):470-479. [doi: [10.1109/TITB.2007.902162](#)] [Medline: [18632327](#)]
38. Pratt W, Unruh K, Civan A, Skeels M. Personal health information management. *Commun. ACM* 2006 Jan 01;49(1):51. [doi: [10.1145/1107458.1107490](#)]
39. Kim SI, Kim HS. Effectiveness of mobile and internet intervention in patients with obese type 2 diabetes. *Int J Med Inform* 2008 Jun;77(6):399-404. [doi: [10.1016/j.ijmedinf.2007.07.006](#)] [Medline: [17881285](#)]
40. Tang PC, Ash JS, Bates DW, Overhage JM, Sands DZ. Personal health records: definitions, benefits, and strategies for overcoming barriers to adoption. *J Am Med Inform Assoc* 2006 Apr;13(2):121-126 [FREE Full text] [doi: [10.1197/jamia.M2025](#)] [Medline: [16357345](#)]
41. Pagliari C, Detmer D, Singleton P. Potential of electronic personal health records. *BMJ* 2007 Aug 18;335(7615):330-333 [FREE Full text] [doi: [10.1136/bmj.39279.482963.AD](#)] [Medline: [17703042](#)]
42. Kahn JS, Aulakh V, Bosworth A. What it takes: characteristics of the ideal personal health record. *Health Aff (Millwood)* 2009;28(2):369-376 [FREE Full text] [doi: [10.1377/hlthaff.28.2.369](#)] [Medline: [19275992](#)]
43. Ball MJ, Smith C, Bakalar RS. Personal health records: empowering consumers. *J Healthc Inf Manag* 2007;21(1):76-86. [Medline: [17299929](#)]
44. Endsley S, Kibbe DC, Linares A, Colorafi K. An introduction to personal health records. *Fam Pract Manag* 2006 May;13(5):57-62 [FREE Full text] [Medline: [16736906](#)]
45. Halamka JD, Mandl KD, Tang PC. Early experiences with personal health records. *J Am Med Inform Assoc* 2008;15(1):1-7 [FREE Full text] [doi: [10.1197/jamia.M2562](#)] [Medline: [17947615](#)]
46. Kaelber DC, Jha AK, Johnston D, Middleton B, Bates DW. A research agenda for personal health records (PHRs). *J Am Med Inform Assoc* 2008;15(6):729-736 [FREE Full text] [doi: [10.1197/jamia.M2547](#)] [Medline: [18756002](#)]
47. Ahern D, Phalen J, Eaton C. The role of eHealth in patient engagement and quality improvement. *eHealth Solutions for Healthcare Disparities* 2008:75-92. [doi: [10.1007/978-0-387-72815-5_9](#)]
48. Grant RW, Wald JS, Schnipper JL, Gandhi TK, Poon EG, Orav EJ, et al. Practice-linked online personal health records for type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med* 2008 Sep 8;168(16):1776-1782. [doi: [10.1001/archinte.168.16.1776](#)] [Medline: [18779465](#)]
49. Hsu CC, Chang HY, Huang MC, Hwang SJ, Yang YC, Lee YS, et al. HbA1c variability is associated with microalbuminuria development in type 2 diabetes: a 7-year prospective cohort study. *Diabetologia* 2012 Dec;55(12):3163-3172. [doi: [10.1007/s00125-012-2700-4](#)] [Medline: [22923064](#)]
50. Penno G, Solini A, Bonora E, Fondelli C, Orsi E, Zerbini G, Renal Insufficiency And Cardiovascular Events Study Group. HbA1c variability as an independent correlate of nephropathy, but not retinopathy, in patients with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian multicenter study. *Diabetes Care* 2013 Aug;36(8):2301-2310. [doi: [10.2337/dc12-2264](#)] [Medline: [23491522](#)]

Abbreviations

- 3G:** third-generation mobile telecommunication
- AADE7:** American Association of Diabetes Educators 7 Self-Care Behaviors
- AES:** Advanced Encryption Standard
- Ajax:** Asynchronous JavaScript XML
- API:** application programming interface
- BMI:** body mass index
- CCD:** Continuity of Care Document
- CDE:** certified diabetes educator
- DMIS:** disease management information system
- DSME:** diabetes self-management education
- HbA1c:** glycosylated hemoglobin
- HIS:** hospital information system
- HL7:** Health Level 7
- HTTPS:** Hypertext Transfer Protocol Secure

IRB: institutional review board
NHI: National Health Insurance
NTUH: National Taiwan University Hospital
PHR: personal health records
SMBG: self-monitoring of blood glucose
SMS: short message service
SOA: service-oriented architecture
T1: entire duration of the telehealthcare program, from September 2011 to February 2013 (18 months)
T1DM: Type 1 diabetes mellitus
T2: initial stage of the telehealthcare program, from September 2011 to February 2012 (6 months)
T2DM: Type 2 diabetes mellitus
T3: middle stage of the telehealthcare program, from March 2012 to August 2012 (6 months)
T4: last stage of the telehealthcare program, from September 2012 to February 2013 (6 months)
XML: Extensible Markup Language

Edited by G Eysenbach; submitted 02.05.13; peer-reviewed by L Diehl, J Gao, Q Li; comments to author 19.06.13; revised version received 13.07.13; accepted 10.11.13; published 09.12.13

Please cite as:

Chen L, Chuang LM, Chang CH, Wang CS, Wang IC, Chung Y, Peng HY, Chen HC, Hsu YL, Lin YS, Chen HJ, Chang TC, Jiang YD, Lee HC, Tan CT, Chang HL, Lai F

Evaluating Self-Management Behaviors of Diabetic Patients in a Telehealthcare Program: Longitudinal Study Over 18 Months
J Med Internet Res 2013;15(12):e266

URL: <http://www.jmir.org/2013/12/e266/>

doi: [10.2196/jmir.2699](https://doi.org/10.2196/jmir.2699)

PMID: [24323283](https://pubmed.ncbi.nlm.nih.gov/24323283/)

©Lichin Chen, Lee-Ming Chuang, Chia-Hsiun Chang, Chiou-Shiang Wang, I-Ching Wang, Yufang Chung, Hui-Yu Peng, Hui-Chuen Chen, Yu-Ling Hsu, Yu-Sheng Lin, Huang-Jen Chen, Tieng-Chun Chang, Yi-Der Jiang, Hung-Chang Lee, Ching-Ting Tan, Hsin-Lu Chang, Feipei Lai. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 09.12.2013. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.