

Original Paper

Digital Transformation of Rheumatology Care in Germany: Cross-Sectional National Survey

Susann May¹, MPH; Robert Darkow², Dr rer med; Johannes Knitza^{3,4}, MHBA, MD, PhD; Katharina Boy¹, BSc; Philipp Klemm⁵, MD; Martin Heinze^{1,6}, MD; Nicolas Vuillerme^{3,7,8}, PhD; Pascal Petit³, PhD; Patricia Steffens-Korbanka⁹, MFA; Heike Kladny¹⁰, MFA; Johannes Hornig⁹, MD; Peer Aries¹¹, MD; Martin Welcker¹⁰, MD; Felix Muehlensiepen^{1,3}, MPH, Dr rer med, PhD

¹Center for Health Services Research, Faculty of Health Sciences Brandenburg, Brandenburg Medical School Theodor Fontane, Rüdersdorf bei Berlin, Germany

²FH JOANNEUM Gesellschaft mbH, Graz, Austria

³AGEIS, Université Grenoble Alpes, Grenoble, France

⁴Institute for Digital Medicine, University Hospital of Giessen and Marburg, Philipps University Marburg, Marburg, Germany

⁵Department of Rheumatology, Immunology, Osteology, Kerckhoff-Klinik GmbH, Bad Nauheim, Germany

⁶University Clinic for Psychiatry and Psychotherapy, Brandenburg Medical School, Immanuel Hospital Rüdersdorf, Rüdersdorf bei Berlin, Germany

⁷Institut Universitaire de France, Paris, France

⁸LabCom Telecom4Health, Orange Labs & Univ. Grenoble Alpes, CNRS, Inria, Grenoble INP-UGA, Grenoble, France

⁹Rheumapraxis an der Hase, Osnabrück, Germany

¹⁰MVZ für Rheumatologie Dr. Martin Welcker GmbH, Planegg, Germany

¹¹Immunologikum Hamburg, Rheumatologie & Klinische Immunologie, Hamburg, Germany

Corresponding Author:

Felix Muehlensiepen, MPH, Dr rer med, PhD
Center for Health Services Research
Faculty of Health Sciences Brandenburg
Brandenburg Medical School Theodor Fontane
Seebad 82/83
Rüdersdorf bei Berlin, 15562
Germany
Phone: 49 1636055860
Email: Felix.Muehlensiepen@mhb-fontane.de

Abstract

Background: In recent years, health care has undergone a rapid and unprecedented digital transformation. In many fields of specialty care, such as rheumatology, this shift is driven by the growing number of patients and limited resources, leading to increased use of digital health technologies (DHTs) to maintain high-quality clinical care. Previous studies examined user acceptance of individual DHTs in rheumatology, such as telemedicine, video consultations, and mHealth. However, it is essential to conduct cross-technology and continuous analyses of user acceptance and DHT use to maximize the benefits for all relevant stakeholders.

Objective: This study aimed to explore the current acceptance, use, and preferences regarding DHTs among patients in rheumatology care in Germany.

Methods: Rheumatology patients from 3 clinics in Germany were surveyed to understand their perspectives on DHTs. The survey included main themes, including acceptance, preferences, COVID-19's impact, potential, and barriers related to DHTs. The data were analyzed using descriptive statistics and correlation analysis.

Results: Out of 337 participants, 53% (179/337) reported using DHTs. Specific technologies included wearables (72/337, 21%), mHealth apps (71/337, 21%), digital therapeutics (32/337, 9%), electronic prescriptions (30/337, 9%), video consultations (15/337, 4%), and at-home blood self-sampling (3/337, 1%). Nearly two-thirds (220/337, 65%) found DHTs useful, and 69% (233/337) held a generally positive attitude toward DHTs. Attitudes shifted positively during the COVID-19 pandemic for 40% (135/337) of participants. Higher education was more prevalent among DHT users (114/179, 63.7%) compared with nonusers (42/151,

27.8%; $P=.02$). The main potential benefits identified were location-independent use (244/337, 72%) and time-independent use (216/337, 64%). Key barriers included insufficient user knowledge (165/337, 49%) and limited information on DHTs (134/337, 40%).

Conclusions: Patient acceptance and use of DHTs in rheumatology is increasing in Germany. A prospective, standardized monitoring of digital transformation in rheumatology care is highly needed.

(*J Med Internet Res* 2025;27:e52601) doi: [10.2196/52601](https://doi.org/10.2196/52601)

KEYWORDS

telemedicine; digital health technologies; rheumatology; COVID-19; questionnaire; telehealth; eHealth; digital health; survey; rheumatism; Germany; Europe; national; use; experience; attitude; opinion; perception; perspective; acceptance; preference; correlation

Introduction

Health care is undergoing a multidisciplinary digital transformation, which refers to “a process that aims to improve an entity by triggering significant changes to its properties through combinations of information, computing, communication, and connectivity technologies” [1].

Within the realm of rheumatology care, a notable surge in digital technologies has transpired in recent years [2,3]. This occurrence stems from the existing global disparity between the increasing number of patients with rheumatic and musculoskeletal disease (RMD) [4] and the static or even decreasing availability of human resources in the field of rheumatology care [5,6].

Consequently, Miloslavsky and Bolster [7] have discerned the amplified use of telemedicine as a prospective remedy to ensure the continuity of rheumatological care in the future [7]. Concurrently, and accelerated by the COVID-19 pandemic, the European Alliance of Associations for Rheumatology (EULAR) has recently published points to consider for remote care in RMD [8].

In Germany, a country struggling with a shrinking and aging population and a shortage of health care professionals, digital health technologies (DHTs) in rheumatology care have proven effective in improving rheumatology care [2,9]. Prominent among the prevailing DHTs are video consultations, sensors, wearables, digital health applications, and digital therapeutics with varied objectives (symptom checkers, ePRO [electronic patient-reported outcome] documentation, or patient education), social media platforms or messenger platforms [10], which can also be combined with home-based self-sampling [11-13].

Past studies have assessed patients' and physicians' acceptance of DHTs in the domain of rheumatology within Germany. These investigations primarily centered on individual digital modalities, such as telemedicine [13,14], mHealth [15], and video consultations [16,17]. Following the onset of the COVID-19 pandemic, the influence of this global crisis on the adoption of DHTs was also scrutinized [3,17]. It remains essential to continuously analyze the digital transformation in rheumatology to maximize the benefits for all relevant stakeholders. In particular, the patient perspective is critical to ensuring that DHTs actually improve health care, meets patient needs, and builds patient trust in the use of DHTs. Thus, the aim of this study was to explore the current acceptance, use,

and preferences regarding DHTs among patients in rheumatology care in Germany.

Methods

Overview

To explore the patients' perspectives and preferences regarding digital health systems in rheumatology care, we recruited rheumatology patients in 3 outpatient clinics. Inclusion criteria were being aged ≥ 18 years, literate in German, having the physical and mental ability to fill out a paper-pencil questionnaire, and having a diagnosis of RMD. The survey was conducted between February 2023 and April 2023. The 3 clinics were sent 125 questionnaires with the request to distribute the questionnaires to eligible patients.

The questionnaire was created based on a literature review and the results of a qualitative study with health care providers (HCPs) and nurses. The questionnaire addressed the following topics: acceptance of DHTs, use of DHTs, preferences for DHTs, the impact of COVID-19 on digital transformation, potential benefits, and barriers regarding DHTs, and sociodemographic characteristics. The survey was pilot-tested with 10 patients to detect necessary formatting and wording changes. Minor revisions were made accordingly. Please refer to [Multimedia Appendix 1](#) for the German version of the questionnaire and [Multimedia Appendix 2](#) for the translated version in English.

Data were analyzed using quantitative descriptive analysis and correlation analysis supported by SPSS (version 23.0; IBM). For correlation analyses, the data were converted into scores, and the Spearman rank correlation coefficient was calculated. The α error distribution was chosen to be 2-sided, as an undirected correlation is assumed to be open to hypotheses. These correlations had a correlation coefficient higher than 0.29.

Ethical Considerations

This study was approved by the ethics committee of Brandenburg Medical School (E-02-20211028). All patients provided informed consent. The rheumatology clinics received an expense allowance of €150 (US \$ 158) for the distribution and return of the questionnaires. The study data are anonymous. The questionnaires were digitized and then stored at the Center for Mental Health at Brandenburg Medical School.

Results

Patient Characteristics

In total, 337 patients with RMD completed the questionnaire. The number of patients rejecting participation was not measured. The study sample's demographics are shown in [Table 1](#). The

mean age was 52.5 (SD 14.2) years; 64% (219/337) were female, 32% (108/337) were male, 1% were (3/337) nonbinary, and 2% (7/337) did not give any information on gender. On average, participants had been receiving rheumatologic treatment for a median 8 years (mean 9.35 years, SD 7.63; range: 1 month-399 months), and the median time with the diagnosis was 8.9 years (mean 10.2 years, SD 8.33; range: 3 months-579 months).

Table 1. Characteristics of participants.

Characteristics	Participants (N=337), n (%)
Sex	
Female	219 (65)
Male	108 (32)
Nonbinary	3 (1)
No data	7 (2)
Diagnosis	
Axial spondyloarthritis	13 (4)
Fibromyalgia	7 (2)
Psoriatic arthritis	40 (12)
Rheumatoid arthritis	182 (54)
Sjögren syndrome	10 (3)
Spondyloarthritis	21 (6)
Systemic lupus erythematosus	9 (3)
Other	39 (12)
No data	16 (5)
Education level	
>10 years of school completed	156 (46)
9-10 years of school completed	163 (48)
Student	2 (1)
No formal schooling	2 (1)
No data	14 (4)
Vocational training	
Apprenticeship	188 (56)
Bachelor's, master's, or magister's degree, diploma	109 (3)
Started an apprenticeship	6 (2)
Without apprenticeship	11 (3)
No data	23 (7)
Place of residence (number of inhabitants)	
Rural region (<5000)	70 (21)
Small city (>5000-20,000)	73 (22)
Medium city (>20,000-100,000)	63 (19)
Large city (>100,000-1,000,000)	28 (8)
Metropolis city (>1,000,000)	93 (28)
No data	10 (3)

Attitudes Toward DHTs

About half of the patients used DHTs overall (179/337, 53%) (Table 2). Overall, most patients (220/337, 65%) rated DHTs as useful, while a third remained neutral in their assessment of usefulness. Almost two-thirds of the patients (233/337, 69%)

were positive or rather positive toward DHTs. Among all participants, 55% (185/337) stated that their attitude toward DHTs had not changed due to the COVID-19 pandemic. However, 40% (135/337) mentioned that their attitude had become more positive.

Table 2. Attitudes toward digital health technologies.

Items	Participants (N=337), n (%)
Do you use digital health technologies?	
Yes	179 (53)
No	151 (45)
No data	7 (2)
Digital health technologies are useful.	
Strongly agree	99 (29)
Agree	121 (36)
Neutral	99 (29)
Disagree	5 (1)
Strongly disagree	7 (2)
No data	6 (2)
How do you rate your attitude towards digital health services?	
Positive	118 (35)
Rather positive	115 (34)
Neutral	77 (23)
Rather negative	17 (5)
Negative	4 (1)
No data	6 (2)
Did you change your attitude due to the COVID-19 pandemic?	
Yes, the attitude has become more positive	135 (40)
Yes, the attitude has become more negative	8 (2)
No	185 (55)
No data	9 (3)

Use of DHTs

Almost half of the patients (143/337, 42%) were already using email for communication with physicians before the COVID-19 pandemic (Table 3). Currently, this value has increased to 58% (195/337). The number of patients using video consultations was very low: 4% (15/337). However, 14% (48/337) stated that

they would use video consultations in the future. We found that 9% (32/337) were using reimbursed digital health applications (German: Digitale Gesundheitsanwendung; DiGA). Overall, 21% (71/337) were using other mobile health apps. Only 1% (3/337) had used blood self-sampling. Almost a quarter (78/337, 23%) stated that they wanted to use electronic prescriptions in the future.

Table 3. Digital health technology use.

	Used before COVID-19, n (%)	I currently use, n (%)	I will use in the future, n (%)	I am not interested, n (%)	I don't use, n (%)	I don't know, n (%)
Video consultation	7 (2)	15 (4)	48 (14)	14 (4)	191 (57)	53 (16)
Digital therapeutics (DiGA ^a)	14 (4)	32 (9)	38 (11)	10 (3)	154 (46)	86 (26)
mHealth apps (not DiGA)	19 (6)	71 (21)	40 (12)	10 (3)	152 (45)	54 (16)
Wearables	19 (6)	72 (21)	21 (6)	19 (6)	175 (52)	43 (13)
Blood self-sampling at home	2 (1)	3 (1)	9 (3)	50 (15)	187 (55)	99 (29)
Electronic prescription	7 (2)	30 (9)	78 (23)	5 (1)	147 (44)	75 (22)

^aDiGA: digital health application.

Potential Benefits of and Barriers to DHTs

Independence in terms of location (244/337, 72%) and time (216/337, 64%) and generally increased flexibility (170/337, 50%) were cited most often by patients as potential benefits of digital health. The detailed results are displayed in [Table 4](#).

The biggest barrier at present is the lack of information. Thus, 49% (165/337) stated that they do not have sufficient knowledge, and 40% (134/337) stated that they are not sufficiently informed about DHTs. A third (118/337, 35%) stated that technical equipment is a barrier to using the tools.

Table 4. Potential benefits and barriers of digital health technologies.

Items	Participants (N=337), n (%)
Potential benefits	
Location independent use	244 (72)
Time-independent use	216 (64)
Detailed documentation of the course of the disease	139 (41)
Cost savings	134 (40)
More options for accessing information, diagnostics, and therapy	125 (37)
Accessibility	85 (25)
More flexibility	170 (50)
Better preparation for the physician-patient consultation	139 (41)
Needs-based care	68 (20)
No potential benefits	32 (9)
Barriers	
Limited information about digital health services	134 (40)
Insufficient evidence of the benefits of the offers	39 (12)
Poor quality of current offers	32 (9)
Gaps in data protection	92 (27)
Lack of user-friendliness	84 (25)
Lack of accessibility	12 (4)
High costs	12 (4)
Lack of technical equipment (eg, poor Internet connection, old end devices)	118 (35)
Lack of knowledge among users	165 (49)
No need because satisfied with the current analogue solutions	57 (17)

Correlation Analyses

While correlation analysis revealed some relationships between content items ([Table 5](#)), no significant correlations between the content items and sociodemographic data were observed.

Table 5. Results of the correlation analyses.

Content items	Spearman ρ	P values (2-tailed)
Current use		
Positive attitude	0.458	<.001
Number of months diagnosed		
Number of months in rheumatology treatment	0.777	<.001
Potential benefits		
Location-independent use		
Detailed documentation of disease progression	0.355	<.001
Accessibility	0.328	<.001
More flexibility	0.331	<.001
No	-0.525	<.001
Detailed documentation of disease progression		
More options for accessing information, diagnostics, and therapy	0.330	<.001
More flexibility	0.336	<.001
Better preparation for the doctor-patient consultation	0.400	<.001
Needs-based care	0.330	<.001
Cost savings		
Accessibility	0.380	<.001
More flexibility	0.320	<.001
Needs-based care	0.302	<.001
More possibilities to access information, diagnostics, therapy		
Accessibility	0.346	<.001
More flexibility	0.405	<.001
Better preparation for the doctor-patient consultation	0.305	<.001
Accessibility		
More flexibility	0.357	<.001
Needs-based care	0.338	<.001
More flexibility		
Needs-based care	0.380	<.001
None	-0.307	<.001
Better preparation for the doctor-patient consultation		
Needs-based care	0.360	<.001
Barriers		
Lack of technical equipment		
Lack of knowledge among users	0.307	<.001

Discussion

To explore the current patient acceptance, use, and preferences for DHTs in rheumatology care, we performed a paper-pencil questionnaire survey among patients with RMD in Germany.

Principal Findings

More than half (179/337, 53%) of 337 patients reported that they use DHTs. Overall, 21% (72/337) used wearables, 21% (71/337) used mHealth apps, 9% (32/337) used digital

therapeutics (DiGA), 9% (30/337) used electronic prescriptions, 4% (15/337) used video consultations, and 1% (3/337) used at-home blood self-sampling. Nearly two-thirds of the patients with RMD (220/337, 65%) rated DHTs as useful. While 69% (233/337) reported a generally positive attitude toward DHTs, about 40% (135/337) mentioned their attitudes became more positive due to the COVID-19 pandemic. The main potential benefits of DHTs reported by the patients were location-independent use (244/337, 72%) and time-independent use (216/337, 64%). The main barriers included insufficient

knowledge among users (165/337, 49%) and limited information about digital health services (134/337, 40%).

Comparison With Previous Work

Our findings are aligned with previous studies that have examined patients' acceptance of DHTs in rheumatology [3,14-19]. Comparing our results with previous findings, clear trends emerge that further underscore the growing acceptance of DHTs among patients with RMD.

In a survey conducted from September 2019 to December 2019 among 766 German patients with RMD [18], only 51% (364/718) of participants indicated familiarity with the term "telemedicine." A mere 30% (210/690) expressed intentions to try telemedicine in the future, and a total of 21% (139/663) expressed a desire for their rheumatologist to offer telemedicine. In this study, conducted from February 2023 to April 2023, that is, after the COVID-19 period, 53% (179/337) of participants reported using DHTs. It should be noted, however, that while the terms telemedicine and DHTs are closely related, they are not synonymous.

In the 2019 survey, 0.3% of patients with RMD reported having experienced a video consultation with a physician. The current results reveal a 4% (15/337) use rate of video consultations, still representing a modest figure, suggesting that video consultations in rheumatology care remain an exceptional practice. These findings corroborate the outcomes observed by Richter et al [17]. In a survey of rheumatologists in Germany, 27% (55/205) reported offering video consultations during COVID-19 lockdown phases, with the frequency of provided video conferences diminishing as the pandemic progressed.

Knitza et al [15] reported that in 2018/2019, most patients with RMD (68%) believed that using medical apps could be beneficial for their own health [15]. However, out of 193 patients, only 8 (4%) were currently using medical apps. In the fall of 2020, Kernder et al [3] explored patient and rheumatologist attitudes toward digital technologies, particularly digital health applications. Even a higher rate of patients (222/299, 74%) and also rheumatologists (98/129, 76%) believed that digital health apps were valuable for managing RMDs. Compared with Knitza et al [15], our results reveal that digital health app use increased notably (71/337, 21%). In our study, a distinction was made between certified and prescribed digital therapeutics (DiGA) and other nonreimbursed digital health apps. Interestingly, patients reported less use of prescribed DiGAs (32/337, 9%) compared with other noncertified mHealth apps (21%). This discrepancy could be attributed to the absence of DiGAs explicitly tailored for RMDs, whereas non-DiGA mHealth apps fill this gap. A first pilot study evaluating DiGA use in rheumatology [20] revealed high patient acceptance and some clinical benefits, yet poor adherence as a major limiting barrier.

According to Kernder et al [3], 38% (112/299) of patients reported a positive change in attitude due to COVID-19, comparable to 40% (135/337) in this study. The most commonly cited advantages of DHAs were their independence from time and place, which were also expressed by participants in our study.

Our study also inquired about the use and acceptance of blood self-sampling, a prospect that has recently gained significance in future rheumatological care [11-13]. However, our results suggest that self-sampling currently remains largely confined to research settings.

In a recent secondary data analysis [21], we demonstrated that, specifically, older patients with RMD residing in rural areas, who could potentially benefit from telemedicine, currently lack the motivation to embrace it and appear to require additional support. However, these relationships were not confirmed by this study. Nevertheless, our data reaffirm the profound relevance of knowledge in the use of digital technologies in rheumatology care. In line with earlier findings [3,14,19,21], our participants identified "lack of knowledge among users" as the main barrier to use. Besides the patient perspective, the viewpoint of HCPs is also pertinent to the implementation of digital technologies. Lack of knowledge among rheumatologists was previously identified as a major barrier to implementing ePROs in routine care [22]. Conveniently, recent surveys also indicated a positive inclination of HCPs toward digital technologies in rheumatology [3,14,16,19].

Limitations

Our study has several limitations. Despite our best efforts, it is possible that our survey did not capture all important emerging technologies relevant to rheumatology. In addition, the terminology used in this survey for DHTs may have caused confusion among participants, potentially leading to information bias. In addition, our survey reached a selected population from 3 outpatient clinics. This also applies to the selection of study participants, as we assume that individuals with a specific interest in digital health were more likely to have completed our questionnaire. A strength of previous surveys [3,16,17,19,22] is that the survey was paper-pen-based (instead of digital) to minimize selection bias.

Finally, it is essential to acknowledge that numerous research groups are focusing on measuring the use and acceptance in the domain of digital rheumatology. However, these studies often use varying approaches and terminologies and refer to different study populations. Therefore, the comparability of data across studies for the purpose of continuous monitoring of the digital transformation in rheumatology remains limited.

Implications

Considering the heterogeneity regarding digital rheumatology surveys and fast transformation, we advocate for a standardized, regular, survey-based monitoring of the digital transformation in rheumatology from the perspectives of both patients with RMD and HCPs. Ideally, this monitoring should be conducted on an international level, including dedicated societies such as EULAR and the Digital Rheumatology Network. Furthermore, the specific digital technologies, their nomenclature, and other questionnaire content should be harmonized through Delphi studies in collaboration with an international expert board involving input from patients.

The results depict high acceptance regarding DHTs, which is currently limited mainly by a lack of knowledge. Dedicated

education for patients with RMD and the treating HCPs is necessary to foster implementation in routine clinical practice.

Conclusions

The digital transformation in rheumatology care in Germany is progressing. Patient acceptance and use is increasing. This

provides hope that, despite the rising burden of disease and stagnating human resources, digital health can continue to ensure high-quality care for patients with RMD in the future. A prospective, standardized monitoring of digital transformation in rheumatology care is highly needed.

Acknowledgments

The authors would like to thank all participants of this study. We also owe special gratitude to the 3 outpatient clinics and their teams that supported this study and distributed the questionnaires.

This study was conducted in cooperation with RheumaDatenRhePort GbR (RHADAR) and uses data from a health care analysis commissioned by AbbVie Deutschland GmbH & Co KG.

Authors' Contributions

FM and SM had full access to all the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. FM, SM, MW, PA, and JH conceptualized and designed the study. FM, SM, MW, PA, JH, PSK, HK, JK, and PK were involved in the acquisition of data. FM, SM, RD, JK, KB, MH, NV, and PP were involved in the analysis and interpretation of data. All the authors were involved in drafting the manuscript and critically revising it for important intellectual content, and they approved the final version of the manuscript to be submitted for publication.

Conflicts of Interest

MW reports support and grants for projects, talks, and discussions by Abbvie, BMS, Boehringer, Galapagos, Gilead, GSK, Hexal, Janssen, Novartis, Pfizer, and UCB. MW is also a member of RHADAR (RheumaDatenRhePort).

Multimedia Appendix 1

Original questionnaire (German).

[\[PDF File \(Adobe PDF File\), 367 KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Questionnaire - English translation.

[\[PDF File \(Adobe PDF File\), 365 KB-Multimedia Appendix 2\]](#)

References

1. Kraus S, Schiavone F, Pluzhnikova A, Invernizzi AC. Digital transformation in healthcare: analyzing the current state-of-research. *J Bus Res*. 2021;123:557-567. [\[FREE Full text\]](#) [doi: [10.1016/j.jbusres.2020.10.030](https://doi.org/10.1016/j.jbusres.2020.10.030)]
2. Mühlensiepen F, Kurkowski S, Krusche M, Mucke J, Prill R, Heinze M, et al. Digital health transition in rheumatology: a qualitative study. *Int J Environ Res Public Health*. 2021;18(5):2636. [\[FREE Full text\]](#) [doi: [10.3390/ijerph18052636](https://doi.org/10.3390/ijerph18052636)] [Medline: [33807952](https://pubmed.ncbi.nlm.nih.gov/33807952/)]
3. Kernder A, Morf H, Klemm P, Vossen D, Haase I, Mucke J, et al. Digital rheumatology in the era of COVID-19: results of a national patient and physician survey. *RMD Open*. 2021;7(1):e001548. [\[FREE Full text\]](#) [doi: [10.1136/rmdopen-2020-001548](https://doi.org/10.1136/rmdopen-2020-001548)] [Medline: [33622673](https://pubmed.ncbi.nlm.nih.gov/33622673/)]
4. Sebbag E, Felten R, Sages F, Sibilia J, Devilliers H, Arnaud L. The world-wide burden of musculoskeletal diseases: a systematic analysis of the World Health Organization Burden of Diseases database. *Ann Rheum Dis*. 2019;78(6):844-848. [doi: [10.1136/annrheumdis-2019-215142](https://doi.org/10.1136/annrheumdis-2019-215142)] [Medline: [30987966](https://pubmed.ncbi.nlm.nih.gov/30987966/)]
5. Al Maini M, Adelowo F, Al Saleh J, Al Weshahi Y, Burmester GR, Cutolo M, et al. The global challenges and opportunities in the practice of rheumatology: white paper by the World Forum on Rheumatic and Musculoskeletal Diseases. *Clin Rheumatol*. 2015;34(5):819-829. [\[FREE Full text\]](#) [doi: [10.1007/s10067-014-2841-6](https://doi.org/10.1007/s10067-014-2841-6)] [Medline: [25501633](https://pubmed.ncbi.nlm.nih.gov/25501633/)]
6. Ward IM, Schmidt TW, Lappan C, Battafarano DF. How critical is tele-medicine to the rheumatology workforce? *Arthritis Care Res (Hoboken)*. 2016;68(10):1387-1389. [\[FREE Full text\]](#) [doi: [10.1002/acr.22853](https://doi.org/10.1002/acr.22853)] [Medline: [26866514](https://pubmed.ncbi.nlm.nih.gov/26866514/)]
7. Miloslavsky EM, Bolster MB. Addressing the rheumatology workforce shortage: a multifaceted approach. *Semin Arthritis Rheum*. 2020;50(4):791-796. [\[FREE Full text\]](#) [doi: [10.1016/j.semarthrit.2020.05.009](https://doi.org/10.1016/j.semarthrit.2020.05.009)] [Medline: [32540672](https://pubmed.ncbi.nlm.nih.gov/32540672/)]
8. de Thurah A, Bosch P, Marques A, Meissner Y, Mukhtyar CB, Knitza J, et al. 2022 EULAR points to consider for remote care in rheumatic and musculoskeletal diseases. *Ann Rheum Dis*. 2022;81(8):1065-1071. [doi: [10.1136/annrheumdis-2022-222341](https://doi.org/10.1136/annrheumdis-2022-222341)] [Medline: [35470160](https://pubmed.ncbi.nlm.nih.gov/35470160/)]

9. Knitza J, Callhoff J, Chehab G, Hueber A, Kiltz U, Kleyer A, et al. [Position paper of the Commission on Digital Rheumatology of the German Society of Rheumatology: tasks, targets and perspectives for a modern rheumatology]. *Z Rheumatol*. 2020;79(6):562-569. [doi: [10.1007/s00393-020-00834-y](https://doi.org/10.1007/s00393-020-00834-y)] [Medline: [32651681](#)]
10. Kataria S, Ravindran V. Digital health: a new dimension in rheumatology patient care. *Rheumatol Int*. 2018;38(11):1949-1957. [doi: [10.1007/s00296-018-4037-x](https://doi.org/10.1007/s00296-018-4037-x)] [Medline: [29713795](#)]
11. Zarbl J, Eimer E, Gigg C, Bendzuck G, Korinth M, Elling-Audersch C, et al. Remote self-collection of capillary blood using upper arm devices for autoantibody analysis in patients with immune-mediated inflammatory rheumatic diseases. *RMD Open*. 2022;8(2):e002641. [FREE Full text] [doi: [10.1136/rmdopen-2022-002641](https://doi.org/10.1136/rmdopen-2022-002641)] [Medline: [36104118](#)]
12. Muehlensiepen F, May S, Zarbl J, Vogt E, Boy K, Heinze M, et al. At-home blood self-sampling in rheumatology: a qualitative study with patients and health care professionals. *BMC Health Serv Res*. 2022;22(1):1470. [FREE Full text] [doi: [10.1186/s12913-022-08787-5](https://doi.org/10.1186/s12913-022-08787-5)] [Medline: [36461025](#)]
13. Mucke J, Knitza J, Muehlensiepen F, Grahammer M, Stenzel R, Simon D, et al. TELERA-asynchronous TELEmedicine for patients with rheumatoid arthritis: study protocol for a prospective, multi-center, randomized controlled trial. *Front Med (Lausanne)*. 2021;8:791715. [FREE Full text] [doi: [10.3389/fmed.2021.791715](https://doi.org/10.3389/fmed.2021.791715)] [Medline: [34966765](#)]
14. Muehlensiepen F, Knitza J, Marquardt W, Engler J, Hueber A, Welcker M. Acceptance of telerheumatology by rheumatologists and general practitioners in Germany: nationwide cross-sectional survey study. *J Med Internet Res*. 2021;23(3):e23742. [FREE Full text] [doi: [10.2196/23742](https://doi.org/10.2196/23742)] [Medline: [33690147](#)]
15. Knitza J, Simon D, Lambrecht A, Raab C, Tascilar K, Hagen M, et al. Mobile health usage, preferences, barriers, and eHealth literacy in rheumatology: patient survey study. *JMIR Mhealth Uhealth*. 2020;8(8):e19661. [FREE Full text] [doi: [10.2196/19661](https://doi.org/10.2196/19661)] [Medline: [32678796](#)]
16. Vossen D, Knitza J, Klemm P, Haase I, Mucke J, Kernder A, et al. [Acceptance of video consultation among patients with inflammatory rheumatic diseases depends on gender and location: results of an online survey among patients and physicians]. *Z Rheumatol*. 2023;82(2):108-113. [FREE Full text] [doi: [10.1007/s00393-021-01052-w](https://doi.org/10.1007/s00393-021-01052-w)] [Medline: [34448915](#)]
17. Richter JG, Chehab G, Reiter J, Aries P, Muehlensiepen F, Welcker M, et al. Evaluation of the use of video consultation in German rheumatology care before and during the COVID-19 pandemic. *Front Med (Lausanne)*. 2022;9:1052055. [FREE Full text] [doi: [10.3389/fmed.2022.1052055](https://doi.org/10.3389/fmed.2022.1052055)] [Medline: [36507506](#)]
18. Muehlensiepen F, Knitza J, Marquardt W, May S, Krusche M, Hueber A, et al. Opportunities and barriers of telemedicine in rheumatology: a participatory, mixed-methods study. *Int J Environ Res Public Health*. 2021;18(24):13127. [FREE Full text] [doi: [10.3390/ijerph182413127](https://doi.org/10.3390/ijerph182413127)] [Medline: [34948737](#)]
19. Becker C, Diener M, Hueber A, Henes J, Krusche M, Ignatyev Y, et al. Unmet information needs of patients with rheumatic diseases: results of a cross-sectional online survey study in Germany. *Int J Environ Res Public Health*. 2022;19(12):7071. [FREE Full text] [doi: [10.3390/ijerph19127071](https://doi.org/10.3390/ijerph19127071)] [Medline: [35742318](#)]
20. Labinsky H, Gupta L, Raimondo MG, Schett G, Knitza J. Real-world usage of digital health applications (DiGA) in rheumatology: results from a German patient survey. *Rheumatol Int*. 2023;43(4):713-719. [FREE Full text] [doi: [10.1007/s00296-022-05261-7](https://doi.org/10.1007/s00296-022-05261-7)] [Medline: [36543961](#)]
21. Muehlensiepen F, Petit P, Knitza J, Welcker M, Vuillerme N. Factors associated with telemedicine use among patients with rheumatic and musculoskeletal disease: secondary analysis of data from a German nationwide survey. *J Med Internet Res*. 2023;25:e40912. [FREE Full text] [doi: [10.2196/40912](https://doi.org/10.2196/40912)] [Medline: [36705950](#)]
22. Krusche M, Klemm P, Grahammer M, Mucke J, Vossen D, Kleyer A, et al. Acceptance, usage, and barriers of electronic patient-reported outcomes among German rheumatologists: survey study. *JMIR Mhealth Uhealth*. 2020;8(7):e18117. [FREE Full text] [doi: [10.2196/18117](https://doi.org/10.2196/18117)] [Medline: [32390592](#)]

Abbreviations

- DHT:** digital health technology
- DiGA:** digital health application
- ePRO:** electronic patient-reported outcome
- EULAR:** European Alliance of Associations for Rheumatology
- HCP:** health care provider
- mHealth:** mobile health
- RMD:** rheumatic and musculoskeletal disease

Edited by N Cahill; submitted 09.09.23; peer-reviewed by M de Zwaan, S Góngora Alonso; comments to author 18.04.24; revised version received 28.05.24; accepted 09.10.24; published 06.01.25

Please cite as:

May S, Darkow R, Knitza J, Boy K, Klemm P, Heinze M, Vuillerme N, Petit P, Steffens-Korbanka P, Kladny H, Hornig J, Aries P, Welcker M, Muehlensiepen F

Digital Transformation of Rheumatology Care in Germany: Cross-Sectional National Survey

J Med Internet Res 2025;27:e52601

URL: <https://www.jmir.org/2025/1/e52601>

doi: [10.2196/52601](https://doi.org/10.2196/52601)

PMID:

©Susann May, Robert Darkow, Johannes Knitza, Katharina Boy, Philipp Klemm, Martin Heinze, Nicolas Vuillerme, Pascal Petit, Patricia Steffens-Korbanka, Heike Kladny, Johannes Hornig, Peer Aries, Martin Welcker, Felix Muehlensiepen. Originally published in the Journal of Medical Internet Research (<https://www.jmir.org>), 06.01.2025. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research (ISSN 1438-8871), is properly cited. The complete bibliographic information, a link to the original publication on <https://www.jmir.org/>, as well as this copyright and license information must be included.