Original Paper

Retracted: Evaluating the Clinical Efficacy of an Exergame-Based Training Program for Enhancing Physical and Cognitive Functions in Older Adults With Mild Cognitive Impairment and Dementia Residing in Rural Long-Term Care Facilities: Randomized Controlled Trial

Aoyu Li¹, PhD; Wei Qiang², PhD; Jingwen Li³, BD; Yan Geng⁴, MD; Yan Qiang⁵, PhD; Juanjuan Zhao¹, PhD

¹School of Software, Taiyuan University of Technology, Jingzhong, China

²Institute of Software, Chinese Academy of Science, Beijing, China

³School of Computer Science, Xijing University, Xi'an, China

⁴Clinical Laboratory and Geriatric, Shanxi Provincial People's Hospital, Taiyuan, China

⁵School of Software, North University of China, Taiyuan, China

Corresponding Author:

Juanjuan Zhao, PhD School of Software Taiyuan University of Technology No. 319, University Street, Yuji District Jingzhong, 030600 China Phone: 86 18636664123 Email: <u>zhaojuanjuan@tyut.edu.cn</u>

Related Articles:

Companion article: <u>https://www.jmir.org/2025/1/e75355/</u> Do not cite this article - this article has been retracted.Retraction notice: <u>https://www.jmir.org/2025/1/e77638</u>

Abstract

Background: Cognitive impairment is an important public health challenge among older adults, particularly in long-term care facilities (LTCFs), where prevalence is higher due to staffing shortages, limited resources, and difficulty maintaining structured exercise programs. Furthermore, older adults often lose interest in repetitive interventions. The exergame "WarioWare: Move It!" (Nintendo) offers a novel solution by combining aerobic exercise, motor coordination, balance training, and cognitive engagement into an immersive experience.

Objective: This study aimed to assess the clinical efficacy of an exergame-based training program delivered via "WarioWare: Move It!" in improving physical flexibility, joint range of motion, motor coordination, hand dexterity, and cognitive function in older adults living in LTCFs.

Methods: The training program was conducted across multiple rural LTCFs in Shanxi Province, China. Participants were randomly assigned to the intervention or control group. The intervention protocol encompassed two 60-minute sessions per week over 12 weeks, using motion-sensing exercises such as waving, jumping, arm swinging, rotational movements, and object-mimicking postures with Joy-Con controllers. Primary outcome measures were derived through clinical tests, including the sit and reach test, shoulder flexibility test, trunk rotation flexibility test, shoulder and elbow range of motion, figure-of-8 walk test, standing balance test, hand dexterity test, and cognitive function tests. Statistical analysis was performed using mixed ANOVA, with time as the within-participant factor and intervention group as the between-participant factor, to assess the training effects on the various outcome measures.

Results: A total of 232 participants were recruited, including 32 (13.8%) patients with mild dementia, 18 (7.8%) with moderate dementia, and 182 (78.4%) with mild cognitive impairment, all of whom completed the study. The mixed ANOVA revealed significant group × time interactions across multiple physical flexibility assessments, including the remaining distance between the hands and toes during the forward bend ($F_{2.156}$ =8.484; P<.001; η^2 =0.098), the distance between the hands clasped behind the

back ($F_{2,156}$ =3.666; P=.04; η^2 =0.045), and the angle formed by trunk rotation to the left and right ($F_{2,156}$ =17.353; P<.001; η^2 =0.182). Significant group × time interactions also emerged for shoulder joint forward flexion ($F_{2,156}$ =17.655; P<.001; η^2 =0.185), abduction ($F_{2,156}$ =6.281; P=.004; η^2 =0.075), and elbow flexion ($F_{2,156}$ =3.298; P=.049; η^2 =0.041). In addition, the time to complete the figure-of-8 walk test ($F_{2,156}$ =11.846; P<.001; η^2 =0.132) and the number of blocks moved within 1 minute ($F_{2,156}$ =4.016; P=.02; η^2 =0.049) showed significant interactions. Finally, all scale-based measures exhibited statistically significant group × time interactions (all P values <.001).

Conclusions: The "WarioWare: Move It!" intervention significantly improved physical flexibility, joint range of motion, motor coordination, hand dexterity, and cognitive function among older adults with mild cognitive impairment or dementia residing in rural LTCFs. The intervention offers an innovative and feasible approach for promoting the health of older adults in resource-limited settings, demonstrating its potential for widespread application in diverse low-resource environments.

Trial Registration: Clinical Trials.gov NCT06717971; https://clinicaltrials.gov/study/NCT06717971

(J Med Internet Res 2025;27:e69109) doi: 10.2196/69109

KEYWORDS

exergame; mild cognitive impairment; dementia; long-term care facilities; multicomponent training

Introduction

Background

Mild cognitive impairment (MCI) and dementia exhibit high prevalence rates among the global older adult population, with new cases projected to increase due to population aging [1,2]. MCI is considered an intermediary stage between age-related cognitive decline and dementia, with studies indicating that approximately 5% to 13% of individuals with MCI progress to dementia each year [3]. The progression is influenced by multiple risk factors, including age, family history, and lifestyle choices, underscoring the critical importance of early intervention and preventive strategies in reducing the incidence of dementia. Dementia, a neurodegenerative disorder with multifactorial etiology, is characterized by the progressive degeneration of brain neurons, resulting in cognitive deficits across several domains, such as complex attention, memory, learning ability, perceptual motor skills, social cognition, and language ability [4]. Due to its multifactorial nature, no effective cure for dementia currently exists, which not only leads to a significant decline in patients' quality of life but also imposes a substantial economic and social burden on families, caregivers, and health care systems [5]. Consequently, early prevention of dementia and maintenance of cognitive function are of particular importance in the older adult population.

In recent years, exergaming, an innovative intervention that integrates physical activity with gaming, has increasingly been introduced into training and rehabilitation programs for the older adult population [6,7]. By combining interactive experiences with immersive virtual environments, exergames encourage active participation among older adults, enhancing cognitive function while alleviating the monotony of traditional exercise forms. Studies indicate that many commercial gaming systems (such as Wii and Kinect) and custom-developed virtual reality games significantly stimulate cognitive abilities, especially in improving memory, attention, and executive function [8,9].

Globally, exergames have garnered significant attention as a promising approach to promoting healthy aging and addressing

```
https://www.jmir.org/2025/1/e69109
```

age-related cognitive and physical decline. Studies conducted across various countries have explored the potential of exergames to combat cognitive impairments, reporting improvements in cognitive and physical health as well as increased enjoyment and motivation among older adults in North America, Europe, and Asia [10-12]. For instance, Kannan et al [13] focused on integrating exergames into community programs and long-term care facilities (LTCFs), demonstrating their effectiveness in enhancing physical functions such as dynamic balance, gait performance, and muscle strength among community-dwelling older adults with reduced physical capacity. Similarly, Chan et al [14] highlighted the role of gamified interventions in fostering social interaction and alleviating feelings of loneliness among older adults, underscoring their cultural adaptability across different social contexts. In Asia, where aging populations are rapidly growing, researchers have developed culturally tailored exergame systems that incorporate traditional activities and movements to enhance their relevance and adherence for older users [15,16].

Moreover, exergames address common barriers to physical activity, such as adverse weather conditions, limited mobility, and geographic isolation, making them particularly valuable in regions with resource constraints or limited access to recreational facilities. These challenges are especially pronounced in rural LTCFs, where cultural attitudes toward exercise often regard physical activity as unnecessary for older adults, and social norms prioritize labor-intensive activities over structured exercise programs. In addition, logistical barriers such as transportation challenges, linguistic diversity, and lack of familiarity with digital technologies hinder the implementation of innovative interventions such as exergames [17,18]. Coupled with the higher prevalence of social isolation in rural settings, these factors necessitate tailored, accessible, and engaging solutions to promote physical and cognitive health among older adults. As global efforts continue to expand the application of exergames in diverse settings, they demonstrate considerable potential for improving the quality of life among older adults, regardless of cultural or socioeconomic contexts.

Despite the promising outcomes of exergaming for older urban populations, its adoption in rural LTCFs remains challenging

XSL•FO RenderX

due to limited health care and social resources [19]. In these settings, shortages of trained caregivers and limited equipment hinder the effective implementation of structured exercise programs, limiting older adults' access to sustained, systematic rehabilitation. In addition, older adults often experience boredom with repetitive exercise routines. They may lack intrinsic motivation, especially when deprived of novel experiences, which can diminish interest and compromise the long-term effectiveness of training [20]. Therefore, exploring and evaluating exergaming systems that are feasible for rural LTCFs and capable of fostering high engagement could serve as a viable approach to enhancing cognitive and physical health among older adults in these communities.

Objectives

Despite the demonstrated clinical benefits of exergames for community-dwelling older adults [6,21], their effectiveness within LTCFs remains insufficiently explored. Current research primarily focuses on improvements in health-related quality of life, cognitive function, and overall functional status [12,22,23], with relatively few assessments targeting specific physical functions, such as flexibility, joint mobility, motor coordination, and hand dexterity. However, these physical capabilities are crucial for the independence of older adults. For instance, limited flexibility can hinder daily activities, directly impacting self-care abilities and increasing dependency on others [24,25]. A reduction in joint mobility is often associated with joint stiffness and pain, restricting the range of motion and diminishing overall physical function, and limiting the living space and activity radius of older adults [26]. Motor coordination is critical in balance maintenance and complex movements, where diminished coordination heightens the risk of falls [27]. Falls can have severe consequences for older adults, including fractures, prolonged immobility, or even accelerated frailty, substantially reducing the quality of life [28-30]. Hand dexterity also underpins the self-sufficiency of older adults. Loss of fine motor skills, such as gripping, pinching, and writing, can restrict their ability to perform daily tasks and participate socially, potentially leading to psychological distress [31].

The Nintendo exergame "WarioWare: Move It!" integrates 12 distinct movement modes, offering not only aerobic, coordination, and balance training but also cognitive exercises that engage memory, attention, and logical reasoning. The game's difficulty can be flexibly adjusted to match users' needs, ensuring optimal training effects at appropriate challenge levels. Despite its evident potential, comprehensive empirical studies are still required to substantiate its benefits fully. Therefore, this study conducted a 12-week longitudinal user study to examine the feasibility and potential clinical utility of "WarioWare: Move It!" for older adult populations in rural LTCFs. Furthermore, this research provides valuable insights and references for older adult rehabilitation professionals and human-computer interaction researchers interested in designing exergame-based interventions for the care of older adults.

Methods

Ethical Considerations

This study was conducted in accordance with the Helsinki Declaration and received approval from the Biomedical Ethics Review Committee of Taiyuan University of Technology (20230641). The study design adheres to the 2013 SPIRIT statement for randomized controlled trials and the CONSORT-EHEALTH (Consolidated Standards of Reporting Trials of Electronic and Mobile Health Applications and Online Telehealth) guidelines (Multimedia Appendix 1), and it has been registered on ClinicalTrials.gov (NCT06717971). Before submitting the written informed consent, participants were informed of potential risks, including minor muscle discomfort or fatigue. They were also made aware that participation was voluntary and that they could withdraw at any time without penalty. All participants provided written informed consent, and eligible participants received a US \$10 compensation.

Participants

In this study, we recruited 232 participants (n=32, 13.8% patients with mild dementia; n=18, 7.8% with moderate dementia; and n=182, 78.4% with MCI) from LTCFs, such as rural daycare centers and nursing homes in Shanxi Province, China. Participant selection followed a purposive sampling method [32], with the entire recruitment process rigorously supervised by experienced neurologists. Caregivers and facility staff were not involved in the participant selection process to avoid potential bias. Inclusion criteria were as follows: (1) normal or corrected-to-normal hearing and vision; (2) age >65 years; (3) residence in an older adult care facility or a minimum stay of 1 month in such a facility; (4) completion of the Global Deterioration Scale (GDS), the Chinese version of the Mini-Mental State Examination (MMSE), and the Chinese version of the Montreal Cognitive Assessment (MoCA), with the capacity to communicate effectively; (5) ability to engage in moderate physical activity without physical disability; (6) absence of severe depressive symptoms or other neurological disorders (eg, stroke, dizziness, and epilepsy); (7) absence of chronic conditions significantly affecting mobility (eg, advanced arthritis, severe osteoporosis, or recent fractures); and (8) provision of informed consent by the participant or their guardian.

All participants underwent comprehensive medical evaluations conducted by experienced neurologists, including a thorough medical history, physical examination, neuroimaging studies (either brain magnetic resonance imaging or computed tomography), and cognitive function assessments. Neurologists assessed participants' cognitive function using the GDS score, classifying the severity of dementia symptoms accordingly. Among the participants with dementia, the average GDS score was 3.7 (SD 0.75), with an average duration of dementia symptoms spanning 3.15 (SD 0.8) years. Cognitive function in the 182 patients with MCI was evaluated using the MoCA, with scores consistently <26. Furthermore, neuroimaging revealed no other structural abnormalities in any participant that could contribute to cognitive impairment. Table 1 summarizes the

```
XSL•FO
```

clinical and demographic characteristics of participants allocated to the intervention and control groups.

Table 1. Clinical and demographic characteristics of participants randomly assigned to the intervention and control groups (N=232).

Characteristics	Intervention group (n=116)	Control group (n=116)	P value
Age (years), mean (SD)	73.03 (4.431)	72.70 (4.958)	.59
Gender, n (%)			.43
Woman	68 (58.6)	62 (53.4)	
Man	48 (41.4)	54 (46.6)	
Hand preference, n (%)			.68
Left	12 (10.3)	14 (12.1)	
Right	104 (89.7)	102 (87.9)	
Exercise habit, n (%)			.43
Yes	62 (53.4)	60 (51.7)	
No	54 (46.6)	56 (48.3)	
Years of education, mean (SD)	6.53 (3.037)	6.31 (2.994)	.57
Hours of sleep, mean (SD)	6.44 (1.239)	6.42 (1.259)	.92
Years of smart device use, mean (SD)	4.59 (1.658)	4.35 (1.741)	.28

Randomization and Blinding

To eliminate selection bias, enhance the internal validity of the research findings, and ensure scientific rigor and reliability, this study used a stringent randomized grouping method to assign participants to either the intervention or control group. The randomization process followed the following strict steps:

- All eligible participants were assigned a unique identification number to maintain anonymity.
- Identification numbers were recorded on slips of paper, which were then sealed in opaque envelopes to ensure a double-anonymized grouping process.
- The numbered envelopes were placed in an opaque container and thoroughly mixed to guarantee complete randomness in selection.
- Researchers wore blindfolds to prevent prior knowledge of envelope contents and randomly drew a single envelope.
- The selection process was repeated until 116 envelopes were chosen to form the intervention group, with the remaining participants allocated to the control group.

After random assignment at a 1:1 ratio, the intervention group received a 12-week training regimen, while the control group continued with standard care provided by LTCFs. Both groups maintained equivalent activity durations.

Intervention Group Training Program

The training program for the intervention group used the Nintendo Switch exergame "WarioWare: Move It," as depicted in Figure 1. Participants engaged with the game through Joy-Con controllers, completing rapid microgames designed to meet specific movement-oriented objectives. Each game session lasted approximately 5 seconds, requiring participants to quickly adapt to various physical actions and postures, such as waving, jumping, swinging arms, rotating, and mimicking object movements. The game's interactive design fosters an engaging experience that promotes physical coordination, reaction speed, upper and lower limb strength, and hand-eye coordination through diverse movement patterns. Switching between tasks compels players to swiftly adjust to varying motion demands, enhancing physical agility while strengthening attention shifting and quick decision-making skills.



Figure 1. Training scenarios and gameplay modes of the Nintendo Switch "WarioWare: Move It" exergame: (A) a participant engaging in a gameplay scenario, (B) competitive mode, (C) cooperative mode, and (D) versus mode.



Furthermore, to enhance both engagement and safety, the intervention program was supported by trained staff responsible for providing tailored assistance to participants throughout the training. Before the intervention, all staff members received comprehensive training in using both the Nintendo Switch console and the game mechanics of "WarioWare: Move It!." The training included familiarization with the game's functionality, understanding how to adapt the tasks to meet participants' physical and cognitive needs, and recognizing when to modify the intensity or duration of tasks. Staff were also trained to monitor participants' physical and cognitive responses, offer guidance as needed, and adjust the game's parameters based on individual performance.

The training program was administered twice weekly, with each session lasting 60 minutes over 12 weeks. Each session comprised three 15-minute gaming segments interspersed with 1- to 5-minute rest periods to sustain participant engagement and mitigate mental fatigue. Before the intervention, all participants attended a 30-minute orientation session to familiarize themselves with the Nintendo Switch console, the game mechanics of "WarioWare: Move It!," and the use of Joy-Con controllers. During training, staff provided tailored support and dynamically adjusted program parameters based

https://www.jmir.org/2025/1/e69109

RenderX

on individual performance, modifying task content, intensity, duration, and physical demands as necessary.

Control Group Standard Care Program

The standard care program for the control group is grounded in routine practices of LTCFs, aiming to sustain and enhance the physical and mental well-being of older adult residents through structured group activities. These activities encompass tailored fitness exercises, horticultural therapy, and sedentary group engagements for older adults. The fitness exercises are adapted to meet the varying physical conditions of participants, ensuring a safe and comfortable environment that promotes moderate physical activity. Horticultural therapy involves hands-on planting and tending to plants, stimulating the senses, and fostering emotional regulation, contributing to improved psychological health. Sedentary activities, such as tabletop games, emphasize social interaction and cognitive maintenance by promoting a relaxed social environment that helps participants remain cognitively engaged. A professional therapist leads each session to ensure that participants receive adequate guidance and support throughout the activities. The program is conducted twice a week, with each session lasting 60 minutes over 12 weeks.

Outcomes Measured

All participants underwent 3 assessments at distinct time points. The initial assessment occurred at baseline before randomization. Subsequent evaluations were conducted at the end of 12 weeks (posttest) and 6 months after the intervention concluded. The primary outcomes of this study focus on detecting intervention effects using indicators related to physical flexibility, joint range of motion, motor coordination, hand dexterity, and cognitive abilities.

In this study, physical flexibility assessments involve several standardized tests. First, the sit and reach test measures the participant's ability to stretch forward while seated with legs extended and heels fixed [33,34]. The farthest distance reached with outstretched hands toward the toes is recorded, with the gap between the fingertips and toes serving as the flexibility index. The shoulder flexibility test assesses flexibility by measuring the distance between the hands when clasped behind the back [35,36]. Participants with higher flexibility achieve either easy hand clasping or shorter hand-to-hand distance. The trunk rotation flexibility test requires participants to stand with feet together and arms extended forward, rotating the torso left and right around the waist [37]. The maximum angle reached on each side indicates flexibility, with more prominent angles signifying greater flexibility.

The joint range of motion reflects the extent of freedom a joint possesses within a specified range of movement. Observing changes in the range of motion across different directions in the intervention group makes it possible to determine whether significant improvements occur in particular directions (eg, flexion or internal rotation), thus providing targeted feedback for the intervention. This study's joint range of motion assessments include shoulder and elbow range of motion tests. The shoulder range of motion test measures the maximum range in 4 directions (eg, flexion, extension, abduction, and adduction) [38,39]. In contrast, the elbow range of motion test assesses the degree of elbow flexion [40].

In addition, this study evaluates motor coordination, defined as the body's ability to synchronize various parts during complex physical tasks. The assessment methods include the figure-of-8 walk [41,42] and standing balance test [43,44]. In the figure-of-8 walk test, participants are instructed to walk in a figure-of-8 pattern around 2 markers placed on the ground, covering a total distance of 10 m. The time taken to complete this task is recorded as a measure of coordination. The standing balance test requires participants to stand with their feet together and their eyes closed, aiming to maintain balance, with the duration of balance maintained being measured as an indicator of stability. Similarly, hand dexterity reflects an individual's coordination, agility, and reaction speed in performing fine motor tasks, particularly when required to complete repetitive, high-precision tasks within a limited time frame. Hand dexterity was assessed using the box block test [45,46], where participants were instructed to use their dominant hand to transfer blocks from one compartment to another within a wooden box over 60 seconds, recording the total number of transfers completed within the specified time. Finally, cognitive function was evaluated using the Cognitive Abilities Screening Instrument (CASI) [47] as well as the Chinese versions of the MMSE [48] and MoCA scales [49]. Detailed measurement procedures are provided in Multimedia Appendix 2.

Statistical Analysis

Statistical analysis was conducted using SPSS for Windows (version 21.0; IBM Corp). Continuous variables were represented by means and SDs, while categorical variables were expressed as frequencies and percentages. Before each analysis, normality and homogeneity of variance were preliminarily assessed. For variables conforming to a normal distribution, we applied 2-tailed *t* tests for continuous variables and chi-square tests for categorical variables to determine the significance of differences between groups. The statistical significance level for all tests was set at P<.05.

To assess the impact of the intervention on outcomes across different time points, we performed a mixed ANOVA, with time as a within- participant factor and the nature of the intervention as a between-participant factor. Effect sizes were calculated using η^2 , which ranges from 0 to 1, to quantify the strength of the intervention's influence. Generally, higher η^2 values indicate a more significant proportion of variance in the dependent variable explained by the intervention, thus reflecting a more substantial intervention effect. For any significant interactions or main effects, Bonferroni correction was used for post hoc evaluations.

Results

Baseline Performance of the Intervention and Control Groups

In this study, a total of 313 participants were screened for eligibility at LTCFs, including rural daycare centers and nursing homes. Of these, 232 participants were included and randomly assigned to either the intervention group (n=116) or the control group (n=116) (Figure 2). Notably, no dropouts occurred during the 12-week intervention period or the 6-month follow-up after the intervention.



Li et al

Figure 2. CONSORT (Consolidated Standards of Reporting Trials) flow diagram.



This study assessed baseline performance for both the intervention and control groups before the intervention began. The evaluation covered multiple domains, including physical flexibility, joint mobility, balance, coordination, and cognitive abilities. As presented in Multimedia Appendix 3, no significant differences were observed between the 2 groups across these various assessments (P>.05), indicating that participants exhibited relatively comparable physical function and cognitive status before the intervention.

Comparison of Physical Function Outcomes Between Intervention and Control Groups

As detailed in Multimedia Appendix 3, the sit and reach test results indicated a significant reduction in the distance between participants' hands and toes over time ($F_{2,78}=35.682$; P<.001). A significant group × time interaction effect was also observed ($F_{2,156}=8.484$; P<.001), demonstrating the intervention group exhibited a markedly more significant flexibility improvement than the control group. A similar trend was observed in the

https://www.jmir.org/2025/1/e69109

RenderX

shoulder flexibility test, where the distance between participants' hands positioned behind the back showed a significant time effect ($F_{2.78}$ =16.511; P<.001) and a significant group × time interaction effect ($F_{2,156}$ =3.666; P=.04). In addition, in the trunk rotation flexibility test, the intervention group exhibited a significant increase in rotational range, with a significant interaction effect $(F_{2,156}=17.353; P<.001)$ and а moderate-to-small effect size ($\eta^2=0.182$). Post hoc analysis demonstrated that physical flexibility indicators in the intervention group showed substantial improvement from baseline to the end of the intervention and remained consistently enhanced up to 6 months post intervention, indicating the durability and stability of the intervention effects.

Furthermore, the joint range of motion was assessed by comparing participants' maximal range of motion in 4 directions (eg, shoulder flexion, extension, abduction, and adduction). The results indicated a significant increase over time in shoulder flexion ($F_{2,78}$ =36.315; P<.001) and abduction ($F_{2,78}$ =4.499; P=.02) among participants. Further analysis revealed a

significant group × time interaction effect, with participants receiving the intervention demonstrating more pronounced improvements in shoulder flexion ($F_{2,156}$ =17.655; P<.001) and abduction ($F_{2,156}$ =6.281, P=.004), suggesting a positive impact of the intervention on enhancing range of motion in specific directions. In addition, the elbow range of motion test showed a significant increase in elbow flexion over time ($F_{2,78}$ =9.261; P<.001), along with a significant group × time interaction effect ($F_{2,156}$ =17.353; P<.001) and a small effect size (η^2 =0.041). These effects may be related to the nature of the intervention tasks (eg, simulated swimming, skiing, and flying), emphasizing shoulder flexion, abduction, and elbow flexion, thereby enhancing the range of motion more substantially in these directions.

Subsequently, in the assessment of motor coordination, the figure-of-8 walk and standing balance test were used to evaluate participants' functional performance based on time metrics. The figure-of-8 walk test measured walking coordination by recording completion time, with shorter times indicating better coordination. The standing balance test assessed static balance ability by recording how participants could maintain balance while standing with their eyes closed, with longer durations indicating better balance. Post hoc analysis revealed a significant time effect in the completion time of the figure-of-8 walk test ($F_{2.78}$ =36.038; P<.001) as well as a significant group × time interaction effect (F_{2,156}=11.846; P<.001), suggesting that the intervention group showed a notably more significant improvement in walking speed and motor coordination than the control group. Similarly, the results from the standing balance test demonstrated a significant increase over time in participants' balance duration with eyes closed ($F_{2,78}$ =9.059; P<.001), indicating a positive impact of the intervention on static balance ability.

Finally, the box block test, used to assess hand dexterity, revealed a significant increase in the number of blocks transferred by participants within 1 minute over time ($F_{2,78}$ =18.648; P<.001). In addition, a significant group × time interaction effect was observed ($F_{2,156}$ =4.016; P=.02), indicating that the intervention group demonstrated a substantially greater improvement in hand dexterity than the control group.

Comparison of Cognitive Function Outcomes Between Intervention and Control Groups

This study assessed cognitive function by examining changes in performance on cognitive scales (eg, CASI, MMSE, and MoCA) between the intervention and control groups, as shown in Multimedia Appendix 3. In the CASI assessment, participants' scores significantly increased over time $(F_{2,78}=21.427; P<.001)$, with a notable group × time interaction effect $(F_{2,156}=13.226; P<.001)$, suggesting a positive impact of the intervention on cognitive function. A similar pattern was observed in both the MMSE and MoCA assessments. For the MMSE, scores significantly improved over time, with a strong time effect $(F_{2,78}=94.552; P<.001)$ and a significant group × time interaction effect $(F_{2,156}=53.244; P<.001)$, supporting the pronounced cognitive improvement within the intervention

https://www.jmir.org/2025/1/e69109

RenderX

group. Likewise, MoCA scores demonstrated a significant time effect ($F_{2,78}$ =52.563; P<.001) and a group × time interaction effect ($F_{2,156}$ =15.882; P<.001), reinforcing the intervention's effectiveness in enhancing various aspects of cognitive function. Notably, the effect sizes (η^2) for the group × time interaction effects across the 3 scales were 0.145 for CASI, 0.406 for MMSE, and 0.169 for MoCA. The relatively larger effect size for MMSE suggests its higher sensitivity in detecting the intervention's impact on cognitive outcomes.

Discussion

Advantages of "WarioWare: Move It!" in Physical and Cognitive Rehabilitation for Older Adults

To our knowledge, this is the first study to evaluate the clinical efficacy of an exergame intervention on a range of functional outcomes, including physical flexibility, joint mobility, motor coordination, hand dexterity, and cognitive status. "WarioWare: Move It!" integrates elements of aerobic exercise, physical coordination, balance training, and cognitive engagement into a motion-based game, with real-time feedback mechanisms to enhance the experience. The immersive setting encourages active participation among older adults, which is particularly valuable in LTCFs in rural areas, where limited medical and social resources make it challenging to implement structured exercise programs. Therefore, we explored the application value of "WarioWare: Move It!" in systematic rehabilitation, focusing on its feasibility in improving older adults' participation and promoting functional recovery. The findings reveal that the older adult participants in LTCFs who received the "WarioWare: Move It!" intervention showed significantly improved flexibility, joint mobility, hand dexterity, and cognitive function compared to a standard care control group. It highlights the potential of "WarioWare: Move It!" as an innovative therapeutic strategy, offering an accessible and well-accepted rehabilitation approach that may help mitigate age-related declines in physical function within long-term care settings.

In addition, this study innovatively incorporates "WarioWare: Move It!" into the design of a motion-based intervention program for older adults. Similar to other exergames, it uses role-playing game mechanics and immersive scene design to foster participant engagement through a playful approach [50,51]. The program offers visual feedback (via screen display) and tactile feedback (through the Joy-Con controller) while incorporating diverse movement postures and randomly varied actions, thus extending beyond traditional exergames limited to fingertip control. Notably, "WarioWare: Move It!" presents a multicomponent training scheme that combines aerobic exercise, body coordination, balance training, and cognitive stimulation to enhance participants' cardiorespiratory endurance, balance, flexibility, and cognitive functions. Prior studies have demonstrated that such multicomponent exergaming provides substantial benefits to physical and cognitive functions in community-dwelling older adults due to its accessibility, cost-effectiveness, and time efficiency, earning broad acceptance [52-54]. Consistent with these findings, this study observed that "WarioWare: Move It!" led to marked improvements in participants' physical abilities (eg, coordination, balance, and

flexibility) and cognitive capacities (eg, executive function, attention, and memory).

One unique aspect of the exergame provided by "WarioWare: Move It!" lies in its ability to automatically set an initial difficulty level at each stage and progressively adjust this based on the player's performance. As players become more familiar with specific game scenarios and movement patterns, the system increases the challenge by gradually increasing the frequency and diversity of the required actions, thus prompting players to achieve efficient coordination between body and mind within a short time frame. Such dynamic adaptability enhances the game's appeal and imposes higher demands on participants' reaction speed, coordination, and focus.

Another fascinating aspect of "WarioWare: Move It!" is its thoughtful inclusion of 3 distinct multiplayer modes, each tailored to meet the preferences of different types of players. In addition to a robust single-player training mode, the game introduces competitive, cooperative, and duel modes to enrich social interactions and enhance player engagement and immersion. First, the competitive mode enables players to compete in specific tasks, with victory determined by scoring or task completion speed, fostering a sense of challenge and achievement. Next, the cooperative mode emphasizes teamwork, encouraging players to complete tasks together, thus strengthening team dynamics and promoting social cohesion. Then, the duel mode focuses on direct player versus player confrontations, offering immediate feedback and creating an intense, engaging battle experience. Overall, these multiplayer interaction modes diversify the game's social and interactive dimensions and offer varied engagement pathways for different player types, thereby boosting player loyalty and satisfaction.

Finally, while this study focused on LTCFs, the intervention's design also makes it feasible for implementation in non-LTCF rural environments. The equipment requirements for "WarioWare: Move It!"—a Joy-Con controller and a screen—are minimal, allowing for adaptation in community centers, individual homes, or other communal spaces. With proper training, family members or community volunteers could facilitate the sessions, alleviating the need for professional caregivers. Moreover, the intervention's engaging and intuitive gameplay makes it accessible even for individuals unfamiliar with gaming technology. Expanding such interventions beyond LTCFs could significantly enhance access to physical and cognitive rehabilitation for older adults in resource-limited settings, promoting healthier aging outcomes across diverse rural populations.

Implications of Multidimensional Improvements in Functional Performance Among Older Adults

In this study, we investigated whether early improvement trends in body coordination and flexibility could be identified in older adults during training. Age-related declines in muscle elasticity and neuromuscular control typically negatively affect coordination and flexibility in older adults [55]. Unlike younger individuals, older adults tend to benefit from training primarily through adaptive changes in the neuromuscular system, improving motor control, postural stability, and coordination, rather than relying solely on muscle hypertrophy or strength

```
https://www.jmir.org/2025/1/e69109
```

XSL•FO

gains [56,57]. Our findings indicated that after the 12-week training period, the remaining distance between the fingertips and toes during the sit and reach test as well as the distance between the hands during the shoulder flexibility test were significantly reduced, indicating a positive effect of the intervention on enhancing both lower and upper limb flexibility. In addition, the trunk rotation angle in both directions significantly increased, further demonstrating the improvement in trunk rotation capacity and overall coordination. Although short-term enhancements in muscle mass may be challenging to detect, neuromuscular adaptations have already positively impacted functional performance in the early stages. In other words, through dynamic contextualized training, older adults experienced enhanced neuromuscular control and motor flexibility, leading to faster, observable improvements in coordination and flexibility.

We also observed a significant improvement in participants' range of motion in specific directions of the shoulder joint, such as flexion and abduction. These movements are crucial for many daily activities, such as dressing and reaching, and their enhancement directly impacts functional performance and independence in older adults' everyday lives [58]. The direction-specific improvement is primarily attributed to the contextualized exercises embedded within the intervention (eg, swimming, flying, and skiing simulations), emphasizing functional relevance and practical application, thereby fostering adaptations in the associated muscle groups and neuromuscular pathways. In addition, the range of motion in elbow joint demonstrated a significant group × time interaction effect in flexion angle, reflecting the rapid adaptability of the neuromuscular system during the short-term intervention. Compared to muscle hypertrophy, older adults are more likely to improve motor control and joint flexibility through neuromuscular adaptations in the short term [25,59], leading to prioritized enhancements in specific directions for the shoulder and elbow joints.

Although the gamified intervention in this study primarily targeted upper limb and trunk movements, a significant increase in walking speed was also observed, indicating the potential benefits of diverse game-based training for gait-related abilities. Walking involves coordinated lower limb and trunk movements, gait control, balance adjustment, and core stability [60,61]. In this study, game interventions (eg, racing, skiing, flying, and archery) not only required upper limb and trunk movements but also challenged balance and center of gravity control through dynamic postures (eg, leaning forward, reaching forward, and lateral shifting). The multifaceted posture control and center of gravity shifting training facilitated improvements in postural stability, balance perception, and neuromuscular coordination, ultimately enhancing gait performance. Moreover, the intervention group significantly reduced completion time for the figure-of-8 walk test, indicating improved stability and turn control during dynamic gait tasks. Similarly, increased duration in the standing balance test reflects improved balance, suggesting that the intervention significantly enhanced participants' neuromuscular control in dynamic and balance tasks. Although the training did not directly target lower limb gait, the center of gravity adjustments, posture transitions, and

full-body coordination exercises embedded within the games significantly improved gait performance.

The intervention program demonstrated significant efficacy in enhancing fine motor control and flexibility of the hands. During the intervention, participants performed hand movement tasks with the Joy-Con controller, such as simulating a tug-of-war, making rock-paper-scissors gestures, and quickly catching a flying disc. These tasks required precise hand adjustments across various contexts, emphasizing independent finger movement and wrist flexibility. In addition, the intervention group showed a notable increase in the number of blocks transferred within 1 minute, alongside a significant group × time interaction effect. This contextualized hand movement training facilitated improvements in fine finger control and wrist flexibility among older adults, with implications for daily tasks requiring fine motor skills, such as grasping objects, handling utensils, and using tools [62].

Finally, this study observed significant improvements in participants' cognitive performance across various assessment tools (eg, CASI, MMSE, and MoCA). Existing research indicates that gamified interventions have a therapeutic impact on the cognitive functions of older adults [63,64]. In this study, the gamified environment provided multidimensional stimulation through visual, auditory, and tactile cues, along with task-based prompts and real-time feedback, consistently presenting participants with spatial and logical challenges. Immersion in such virtual scenarios contributed not only to attentional recovery and stress reduction but also played a positive role in cognitive rehabilitation and enhancement [65]. Moreover, the nature of gamified training, which requires participants to make quick decisions in dynamic scenarios, stimulated executive functioning and response capabilities [66]. Notably, the design features of "WarioWare: Move It!" integrated aerobic exercise, coordination, and balance training alongside cognitive tasks involving memory, attention, and logical reasoning. These multilayered cognitive stimulations created a comprehensive training environment, promoting broad activation of brain functions. Therefore, the notable improvements observed in brain health assessments postintervention were anticipated, highlighting the significant role of gamified interventions in cognitive rehabilitation and enhancement for older adults.

Insights for Rehabilitation Professionals and Human-Computer Interaction Researchers in Older Adult Care

This study offers a new perspective for rehabilitation professionals working with older adult populations, focusing on three key areas: (1) integrating multicomponent cognitive training with physical function restoration—"WarioWare: Move It!" effectively combines aerobic exercise, body coordination, balance, and cognitive stimulation within a structured platform that facilitates multifunctional training through various interactive physical movements—and (2) enhancing adherence through practicality and enjoyment—by integrating intuitive visual feedback, goal achievement mechanisms, immersive scenarios, real-time feedback, and progressively challenging tasks, WarioWare: Move It! Fosters a sense of accomplishment and interactive enjoyment, reinforcing experiential learning and

https://www.jmir.org/2025/1/e69109

 $XSI \bullet FC$

significantly improving adherence—and (3) Practical application of targeted physical function training—through contextualized and interactive movement design, "WarioWare: Move It!" incorporates actions frequently performed in daily life (eg, shoulder flexion and abduction, gait balance, and hand movements) within gameplay to enhance specific physical functions (eg, joint mobility and flexibility).

This study offers valuable insights for researchers in the human-computer interaction field, specifically in designing older adult-friendly interactive systems. Specifically, it highlights three key areas: (1) dynamic adaptive design for older adult-friendly interventions-the dynamic adaptability of the system in "WarioWare: Move It!" gradually guides participants toward higher levels of physical and cognitive demands, facilitating synchronized improvements in both domains within a safe and comfortable experience; (2) multisensory feedback and enhanced interaction-the Intervention successfully integrates multisensory stimulation through visual, auditory, and tactile feedback, enhancing the immersive experience and improving the brain's responsiveness through multimodal input; and (3) rehabilitation benefits of multiple social interaction modes-the social interaction modes within the exergames, including competition (which stimulates participants' achievement motivation), cooperation (which strengthens teamwork awareness), and versus challenges (which foster quick decision-making and response skills), not only enrich the social experience but also meet the psychological and emotional needs of older adults, thus enhancing rehabilitation outcomes.

Limitations and Future Work

Our study has several limitations. First, although the sample size exceeds the requirements for statistical analysis, it remains relatively small. Second, although the exergames have positively affected specific physical and cognitive functions, they cannot fully replace traditional rehabilitation methods (eg, physical or cognitive behavioral therapy). Future intervention designs could consider integrating exergames with conventional rehabilitation techniques, creating a more diversified and comprehensive rehabilitation plan to meet the broader needs of older adults. Third, the social interaction features of "WarioWare: Move It!" (eg, competition, cooperation, and rivalry) enhance rehabilitation outcomes. However, preferences for competition or collaborative modes may vary among older adults, particularly those who are more introverted or have lower social engagement, making these modes less appealing to some individuals. Future game iterations could incorporate personalized social interaction options to ensure each participant receives an optimal experience, boosting engagement. Fourth, although this study focused on older adults, differences in age, stage of cognitive impairment, and the specific needs of different age groups may warrant further investigation to understand the broader applicability of the intervention.

Conclusions

The "WarioWare: Move It!" training program significantly improves physical flexibility, joint range of motion, motor coordination, hand dexterity, and cognitive function among older adults in rural LTCFs. For patients with MCI or dementia, this intervention not only aids in delaying functional decline

but also enhances cognitive function and quality of daily living through multisensory stimulation and dynamic physical engagement. Furthermore, the incorporation of exergames marks an innovative shift in older adult care models, offering a novel, low-demand intervention method that effectively addresses the shortcomings of traditional care approaches. These findings support the integration of exergames into routine care practices and provide a viable, technology-driven solution to address the challenges of aging health in institutional care settings worldwide.

Acknowledgments

This research received funding from the National Natural Science Foundation of China (62476190, 62376183, and U21A20469), the Central Government–Guided Science and Technology Development Program (YDZJSX2022C004), and Shanxi Provincial Administration of Traditional Chinese Medicine Scientific Research Project Plan (2024ZYY2A033). The authors sincerely thank the volunteers whose invaluable participation made the successful completion of this study possible.

Data Availability

The datasets generated and analyzed during this study are available from the corresponding author upon reasonable request.

Conflicts of Interest

None declared.

Multimedia Appendix 1

CONSORT-EHEALTH (Consolidated Standards of Reporting Trials of Electronic and Mobile Health Applications and Online Telehealth) checklist.

[PDF File (Adobe PDF File), 359 KB-Multimedia Appendix 1]

Multimedia Appendix 2

An overview of the broad scope of tests and key assessment areas. [PDF File (Adobe PDF File), 110 KB-Multimedia Appendix 2]

Multimedia Appendix 3

Data of measurement outcomes at baseline, posttest, and after 6 months postintervention. [PDF File (Adobe PDF File), 243 KB-Multimedia Appendix 3]

References

- 1. Gauthier S, Reisberg B, Zaudig M, Petersen RC, Ritchie K, Broich K, et al. Mild cognitive impairment. Lancet. Apr 15, 2006;367(9518):1262-1270. [doi: 10.1016/S0140-6736(06)68542-5] [Medline: 16631882]
- GBD 2019 Dementia Forecasting Collaborators. Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. Lancet Public Health. Feb 2022;7(2):e105-e125. [FREE Full text] [doi: 10.1016/S2468-2667(21)00249-8] [Medline: 34998485]
- 3. Petersen RC. Mild cognitive impairment. Continuum (Minneap Minn). Apr 2016;22(2 Dementia):404-418. [FREE Full text] [doi: 10.1212/CON.0000000000313] [Medline: 27042901]
- Sachdev PS, Blacker D, Blazer DG, Ganguli M, Jeste DV, Paulsen JS, et al. Classifying neurocognitive disorders: the DSM-5 approach. Nat Rev Neurol. Nov 30, 2014;10(11):634-642. [FREE Full text] [doi: 10.1038/nrneurol.2014.181] [Medline: 25266297]
- Yiannopoulou KG, Papageorgiou SG. Current and future treatments in Alzheimer disease: an update. J Cent Nerv Syst Dis. Feb 29, 2020;12:1179573520907397. [FREE Full text] [doi: 10.1177/1179573520907397] [Medline: 32165850]
- Pacheco TB, de Medeiros CS, de Oliveira VH, Vieira ER, de Cavalcanti FA. Effectiveness of exergames for improving mobility and balance in older adults: a systematic review and meta-analysis. Syst Rev. Jul 18, 2020;9(1):163. [FREE Full text] [doi: 10.1186/s13643-020-01421-7] [Medline: 32682439]
- Wu S, Ji H, Won J, Jo E, Kim Y, Park J. The effects of exergaming on executive and physical functions in older adults with dementia: randomized controlled trial. J Med Internet Res. Mar 07, 2023;25:e39993. [FREE Full text] [doi: 10.2196/39993] [Medline: <u>36881445</u>]
- Hughes TF, Flatt JD, Fu B, Butters MA, Chang CH, Ganguli M. Interactive video gaming compared with health education in older adults with mild cognitive impairment: a feasibility study. Int J Geriatr Psychiatry. Sep 22, 2014;29(9):890-898.
 [FREE Full text] [doi: 10.1002/gps.4075] [Medline: 24452845]

- Li et al
- 9. Li A, Li J, Wu W, Zhao J, Qiang Y. Effect of virtual reality training on cognitive function and motor performance in older adults with cognitive impairment receiving health care: a randomized controlled trial. Int J Hum Comput Interact. Oct 19, 2023;40(22):7755-7772. [doi: 10.1080/10447318.2023.2271240]
- 10. Karssemeijer EG, Aaronson JA, Bossers WJ, Donders R, Olde Rikkert MG, Kessels RP. The quest for synergy between physical exercise and cognitive stimulation via exergaming in people with dementia: a randomized controlled trial. Alzheimers Res Ther. Jan 05, 2019;11(1):3. [FREE Full text] [doi: 10.1186/s13195-018-0454-z] [Medline: 30611286]
- 11. Kamnardsiri T, Kumfu S, Munkhetvit P, Boripuntakul S, Sungkarat S. Home-based, low-intensity, gamification-based, interactive physical-cognitive training for older adults using the ADDIE model: design, development, and evaluation of user experience. JMIR Serious Games. Oct 29, 2024;12:e59141. [FREE Full text] [doi: 10.2196/59141] [Medline: 39470391]
- Chu CH, Biss RK, Cooper L, Quan AM, Matulis H. Exergaming platform for older adults residing in long-term care homes: user-centered design, development, and usability study. JMIR Serious Games. Mar 09, 2021;9(1):e22370. [FREE Full text] [doi: 10.2196/22370] [Medline: <u>33687337</u>]
- Kannan L, Sahu U, Subramaniam S, Mehta N, Kaur T, Hughes S, et al. Gaming-based tele-exercise program to improve physical function in frail older adults: feasibility randomized controlled trial. J Med Internet Res. Nov 27, 2024;26:e56810. [FREE Full text] [doi: 10.2196/56810] [Medline: 39602215]
- Chan G, Banire B, Anukem S, Imran M, Meena S, Nwagu C, et al. Social exergames in health and wellness: a systematic review of trends, effectiveness, challenges, and directions for future research. Int J Hum Comput Interact. Jul 08, 2024:1-32. [doi: 10.1080/10447318.2024.2371686]
- Du Q, Song Z, Jiang H, Wei X, Weng D, Fan M. LightSword: a customized virtual reality exergame for long-term cognitive inhibition training in older adults. In: Proceedings of the 2024 CHI Conference on Human Factors in Computing Systems. 2024. Presented at: CHI '24; May 11-16, 2024:1-17; Honolulu, HI. URL: <u>https://dl.acm.org/doi/10.1145/3613904.3642187</u> [doi: <u>10.1145/3613904.3642187</u>]
- Chen Q, Wu Z, Liu Y, Han L, Li Z, Kan GL, et al. SilverCycling: exploring the impact of bike-based locomotion on spatial orientation for older adults in VR. Proc ACM Interact Mob Wearable Ubiquitous Technol. Sep 09, 2024;8(3):1-24. [doi: 10.1145/3678522]
- 17. Meekes W, Stanmore EK. Motivational determinants of exergame participation for older people in assisted living facilities: mixed-methods study. J Med Internet Res. Jul 06, 2017;19(7):e238. [FREE Full text] [doi: 10.2196/jmir.6841] [Medline: 28684385]
- Unbehaun D, Vaziri DD, Aal K, Wieching R, Tolmie P, Wulf V. Exploring the potential of exergames to affect the social and daily life of people with dementia and their caregivers. In: Proceedings of the 2018 CHI Conference on Human Factors in Computing Systems. 2018. Presented at: CHI '18; April 21-26, 2018:1-15; Montreal, QC. URL: <u>https://dl.acm.org/doi/ 10.1145/3173574.3173636</u> [doi: 10.1145/3173574.3173636]
- Chen KM, Li CH, Huang HT, Cheng Y. Feasible modalities and long-term effects of elastic band exercises in nursing home older adults in wheelchairs: a cluster randomized controlled trial. Int J Nurs Stud. Mar 2016;55:4-14. [doi: 10.1016/j.ijnurstu.2015.11.004] [Medline: 26655368]
- Karaosmanoglu S, Rings S, Kruse L, Stein C, Steinicke F. Lessons learned from a human-centered design of an immersive exergame for people with dementia. Proc ACM Hum Comput Interact. Oct 06, 2021;5(CHI PLAY):1-27. [doi: 10.1145/3474679]
- 21. Kappen DL, Mirza-Babaei P, Nacke LE. Older adults' physical activity and exergames: a systematic review. Int J Hum Comput Interact. Feb 23, 2018;35(2):140-167. [doi: <u>10.1080/10447318.2018.1441253</u>]
- 22. Wang RY, Huang YC, Zhou JH, Cheng S, Yang Y. Effects of exergame-based dual-task training on executive function and dual-task performance in community-dwelling older people: a randomized-controlled trial. Games Health J. Oct 01, 2021;10(5):347-354. [doi: 10.1089/g4h.2021.0057] [Medline: 34491113]
- 23. Chiu HM, Hsu MC, Ouyang WC. Effects of incorporating virtual reality training intervention into health care on cognitive function and wellbeing in older adults with cognitive impairment: a randomized controlled trial. Int J Hum Comput Interact. Feb 2023;170:102957. [doi: 10.1016/j.ijhcs.2022.102957]
- 24. Seco J, Abecia LC, Echevarría E, Barbero I, Torres-Unda J, Rodriguez V, et al. A long-term physical activity training program increases strength and flexibility, and improves balance in older adults. Rehabil Nurs. 2013;38(1):37-47. [doi: 10.1002/rnj.64] [Medline: 23365004]
- 25. Stathokostas L, Little RM, Vandervoort AA, Paterson DH. Flexibility training and functional ability in older adults: a systematic review. J Aging Res. 2012;2012:306818-306830. [FREE Full text] [doi: 10.1155/2012/306818] [Medline: 23209904]
- 26. Coleman EA, Buchner DM, Cress ME, Chan BK, de Lateur BJ. The relationship of joint symptoms with exercise performance in older adults. J Am Geriatr Soc. Jan 27, 1996;44(1):14-21. [doi: 10.1111/j.1532-5415.1996.tb05632.x] [Medline: 8537585]
- 27. Brach JS, Perera S, Shuman V, Gil AB, Kriska A, Nadkarni NK, et al. Effect of timing and coordination training on mobility and physical activity among community-dwelling older adults: a randomized clinical trial. JAMA Netw Open. May 02, 2022;5(5):e2212921. [FREE Full text] [doi: 10.1001/jamanetworkopen.2022.12921] [Medline: 35604689]
- 28. Ambrose AF, Paul G, Hausdorff JM. Risk factors for falls among older adults: a review of the literature. Maturitas. May 2013;75(1):51-61. [doi: 10.1016/j.maturitas.2013.02.009] [Medline: 23523272]

```
https://www.jmir.org/2025/1/e69109
```

29.

- Chang JT, Morton SC, Rubenstein LZ, Mojica WA, Maglione M, Suttorp MJ, et al. Interventions for the prevention of falls in older adults: systematic review and meta-analysis of randomised clinical trials. BMJ. Mar 20, 2004;328(7441):680. [FREE Full text] [doi: 10.1136/bmj.328.7441.680] [Medline: 15031239]
- Tricco AC, Thomas SM, Veroniki AA, Hamid JS, Cogo E, Strifler L, et al. Comparisons of interventions for preventing falls in older adults: a systematic review and meta-analysis. JAMA. Nov 07, 2017;318(17):1687-1699. [FREE Full text] [doi: 10.1001/jama.2017.15006] [Medline: 29114830]
- Scherder E, Dekker W, Eggermont L. Higher-level hand motor function in aging and (preclinical) dementia: its relationship with (instrumental) activities of daily life--a mini-review. Gerontology. Nov 11, 2008;54(6):333-341. [FREE Full text] [doi: 10.1159/000168203] [Medline: 18997468]
- 32. Polit DF, Beck CT. Nursing Research: Generating and Assessing Evidence for Nursing Practice. 10th edition. New York, NY. Lippincott Williams & Wilkins; 2008.
- 33. Wells KF, Dillon EK. The sit and reach—a test of back and leg flexibility. Res Q Am Assoc Health Phys Educ. Dec 26, 2013;23(1):115-118. [doi: 10.1080/10671188.1952.10761965]
- 34. Jones CJ, Rikli RE, Max J, Noffal G. The reliability and validity of a chair sit-and-reach test as a measure of hamstring flexibility in older adults. Res Q Exerc Sport. Dec 1998;69(4):338-343. [doi: 10.1080/02701367.1998.10607708] [Medline: 9864752]
- 35. Borstad JD, Mathiowetz KM, Minday LE, Prabhu B, Christopherson DE, Ludewig PM. Clinical measurement of posterior shoulder flexibility. Man Ther. Nov 2007;12(4):386-389. [doi: <u>10.1016/j.math.2006.07.014</u>] [Medline: <u>16971161</u>]
- 36. Bakhsh W, Nicandri G. Anatomy and physical examination of the Shoulder. Sports Med Arthrosc Rev. Sep 2018;26(3):e10-e22. [doi: 10.1097/JSA.00000000000202] [Medline: 30059442]
- Merritt JL, McLean TJ, Erickson RP, Offord KP. Measurement of trunk flexibility in normal subjects: reproducibility of three clinical methods. Mayo Clin Proc. Mar 1986;61(3):192-197. [doi: <u>10.1016/s0025-6196(12)61848-5</u>] [Medline: <u>3945120</u>]
- 38. Hayes K, Walton JR, Szomor ZL, Murrell GA. Reliability of five methods for assessing shoulder range of motion. Aust J Physiother. 2001;47(4):289-294. [doi: 10.1016/s0004-9514(14)60274-9] [Medline: 11722295]
- Mullaney MJ, McHugh MP, Johnson CP, Tyler TF. Reliability of shoulder range of motion comparing a goniometer to a digital level. Physiother Theory Pract. Jul 17, 2010;26(5):327-333. [doi: <u>10.3109/09593980903094230</u>] [Medline: <u>20557263</u>]
- 40. Armstrong AD, MacDermid JC, Chinchalkar S, Stevens RS, King GJ. Reliability of range-of-motion measurement in the elbow and forearm. J Shoulder Elbow Surg. Nov 1998;7(6):573-580. [doi: <u>10.1016/s1058-2746(98)90003-9</u>] [Medline: <u>9883416</u>]
- 41. Wong SS, Yam MS, Ng SS. The Figure-of-Eight Walk test: reliability and associations with stroke-specific impairments. Disabil Rehabil. Apr 19, 2013;35(22):1896-1902. [doi: 10.3109/09638288.2013.766274] [Medline: 23600714]
- 42. Dhillon RK, Sharma M. A review on figure of eight walking in physical therapy rehabilitation of elderly. Work Older People. Aug 17, 2021;25(3):214-219. [doi: 10.1108/wwop-04-2021-0020]
- 43. Browne JE, O'Hare NJ. Review of the different methods for assessing standing balance. Physiotherapy. Sep 2001;87(9):489-495. [doi: 10.1016/S0031-9406(05)60696-7]
- 44. Newton R. Review of tests of standing balance abilities. Brain Inj. Jul 03, 1989;3(4):335-343. [doi: 10.3109/02699058909004558] [Medline: 2684312]
- 45. Mathiowetz V, Volland G, Kashman N, Weber K. Adult norms for the Box and Block Test of manual dexterity. Am J Occup Ther. Jun 1985;39(6):386-391. [doi: <u>10.5014/ajot.39.6.386</u>] [Medline: <u>3160243</u>]
- 46. Desrosiers J, Bravo G, Hébert R, Dutil E, Mercier L. Validation of the Box and Block Test as a measure of dexterity of elderly people: reliability, validity, and norms studies. Arch Phys Med Rehabil. Jul 1994;75(7):751-755. [Medline: <u>8024419</u>]
- 47. Teng EL, Hasegawa K, Homma A, Imai Y, Larson E, Graves A, et al. The Cognitive Abilities Screening Instrument (CASI): a practical test for cross-cultural epidemiological studies of dementia. Int Psychogeriatr. Mar 1994;6(1):45-62. [doi: 10.1017/s1041610294001602] [Medline: 8054493]
- 48. Guo NW, Liu HC, Wong PF, Liao KK, Yan SH, Lin KP, et al. Chinese version and norms of the mini-mental state examination. Rehabil Pract Sci. Dec 1, 1988;16(1):52. [doi: <u>10.6315/3005-3846.1745</u>]
- Yu J, Li J, Huang X. The Beijing version of the Montreal Cognitive Assessment as a brief screening tool for mild cognitive impairment: a community-based study. BMC Psychiatry. Sep 25, 2012;12(1):156. [FREE Full text] [doi: 10.1186/1471-244X-12-156] [Medline: 23009126]
- Schättin A, Pickles J, Flagmeier D, Schärer B, Riederer Y, Niedecken S, et al. Development of a novel home-based exergame with on-body feedback: usability study. JMIR Serious Games. Dec 06, 2022;10(4):e38703. [FREE Full text] [doi: 10.2196/38703] [Medline: 36472900]
- Richards C, Graham TC. Developing compelling repetitive-motion exergames by balancing player agency with the constraints of exercise. In: Proceedings of the 2016 ACM Conference on Designing Interactive Systems. 2016. Presented at: DIS '16; June 4-8, 2016:911-923; Brisbane, Australia. URL: <u>https://dl.acm.org/doi/10.1145/2901790.2901824</u> [doi: <u>10.1145/2901790.2901824</u>]

- 52. Moreira NB, Rodacki AL, Costa SN, Pitta A, Bento PC. Perceptive-cognitive and physical function in prefrail older adults: exergaming versus traditional multicomponent training. Rejuvenation Res. Feb 01, 2021;24(1):28-36. [doi: 10.1089/rej.2020.2302] [Medline: 32443963]
- Zhao Y, Feng H, Wu X, Du Y, Yang X, Hu M, et al. Effectiveness of exergaming in improving cognitive and physical function in people with mild cognitive impairment or dementia: systematic review. JMIR Serious Games. Jun 30, 2020;8(2):e16841. [FREE Full text] [doi: 10.2196/16841] [Medline: 32602841]
- Liao YY, Chen IH, Hsu WC, Tseng H, Wang R. Effect of exergaming versus combined exercise on cognitive function and brain activation in frail older adults: a randomised controlled trial. Ann Phys Rehabil Med. Sep 2021;64(5):101492. [FREE Full text] [doi: 10.1016/j.rehab.2021.101492] [Medline: 33454398]
- 55. Bellew JW, Symons BT, Vandervoort AA. Geriatric fitness: effects of aging and recommendations for exercise in older adults. Cardiopulm Phys Ther J. 2005;16(1):20-31. [doi: 10.1097/01823246-200516010-00005]
- 56. Moritani T, deVries HA. Potential for gross muscle hypertrophy in older men. J Gerontol. Sep 01, 1980;35(5):672-682. [doi: <u>10.1093/geronj/35.5.672</u>] [Medline: <u>7430562</u>]
- 57. Macaluso A, De Vito G. Muscle strength, power and adaptations to resistance training in older people. Eur J Appl Physiol. Apr 25, 2004;91(4):450-472. [doi: 10.1007/s00421-003-0991-3] [Medline: 14639481]
- Oosterwijk AM, Nieuwenhuis MK, van der Schans CP, Mouton L. Shoulder and elbow range of motion for the performance of activities of daily living: a systematic review. Physiother Theory Pract. Jul 29, 2018;34(7):505-528. [FREE Full text] [doi: 10.1080/09593985.2017.1422206] [Medline: 29377745]
- 59. Holland GJ, Tanaka K, Shigematsu R, Nakagaichi M. Flexibility and physical functions of older adults: a review. J Aging Phys Act. 2002;10(2):169-206. [FREE Full text] [doi: 10.1123/japa.10.2.169]
- 60. Clark DJ. Automaticity of walking: functional significance, mechanisms, measurement and rehabilitation strategies. Front Hum Neurosci. May 05, 2015;9:246. [FREE Full text] [doi: 10.3389/fnhum.2015.00246] [Medline: 25999838]
- 61. Lacquaniti F, Grasso R, Zago M. Motor patterns in walking. News Physiol Sci. Aug 1999;14(4):168-174. [doi: 10.1152/physiologyonline.1999.14.4.168] [Medline: 11390844]
- 62. Liu B, Chen X, Li Y, Liu H, Guo S, Yu P. Effect of passive finger exercises on grip strength and the ability to perform activities of daily living for older people with dementia: a 12-week randomized controlled trial. Clin Interv Aging. 2018;13:2169-2177. [FREE Full text] [doi: 10.2147/CIA.S174756] [Medline: 30464426]
- 63. Vermeir JF, White MJ, Johnson D, Crombez G, Van Ryckeghem DM. The effects of gamification on computerized cognitive training: systematic review and meta-analysis. JMIR Serious Games. Aug 10, 2020;8(3):e18644. [FREE Full text] [doi: 10.2196/18644] [Medline: 32773374]
- 64. Lumsden J, Edwards EA, Lawrence NS, Coyle D, Munafò MR. Gamification of cognitive assessment and cognitive training: a systematic review of applications and efficacy. JMIR Serious Games. Jul 15, 2016;4(2):e11. [FREE Full text] [doi: 10.2196/games.5888] [Medline: 27421244]
- 65. Yen HY, Chiu HL. Virtual reality exergames for improving older adults' cognition and depression: a systematic review and meta-analysis of randomized control trials. J Am Med Dir Assoc. May 2021;22(5):995-1002. [doi: 10.1016/j.jamda.2021.03.009] [Medline: 33812843]
- 66. Liao YY, Chen IH, Lin YJ, Chen Y, Hsu W. Effects of virtual reality-based physical and cognitive training on executive function and dual-task gait performance in older adults with mild cognitive impairment: a randomized control trial. Front Aging Neurosci. Jul 16, 2019;11:162. [FREE Full text] [doi: 10.3389/fnagi.2019.00162] [Medline: 31379553]

Abbreviations

CASI: Cognitive Abilities Screening Instrument CONSORT-EHEALTH: Consolidated Standards of Reporting Trials of Electronic and Mobile Health Applications and Online Telehealth GDS: Global Deterioration Scale LTCF: long-term care facility MCI: mild cognitive impairment MMSE: Mini-Mental State Examination MoCA: Montreal Cognitive Assessment



Edited by A Mavragani; submitted 22.11.24; peer-reviewed by Y Huusi, T Ramsey, L Yao; comments to author 22.12.24; revised version received 30.12.24; accepted 07.01.25; published 19.02.25 <u>Please cite as:</u> Li A, Qiang W, Li J, Geng Y, Qiang Y, Zhao J Retracted: Evaluating the Clinical Efficacy of an Exergame-Based Training Program for Enhancing Physical and Cognitive Functions in Older Adults With Mild Cognitive Impairment and Dementia Residing in Rural Long-Term Care Facilities: Randomized Controlled Trial J Med Internet Res 2025;27:e69109 URL: https://www.jmir.org/2025/1/e69109 doi: 10.2196/69109 PMID: 39969990

©Aoyu Li, Wei Qiang, Jingwen Li, Yan Geng, Yan Qiang, Juanjuan Zhao. Originally published in the Journal of Medical Internet Research (https://www.jmir.org), 19.02.2025. This is an open-access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research (ISSN 1438-8871), is properly cited. The complete bibliographic information, a link to the original publication on https://www.jmir.org/, as well as this copyright and license information must be included.

