Letter to the Editor

Enhancing Digital Health Interventions for Medication Adherence: Considerations for Broader Applicability and Long-Term Impact

ShanShan Du¹, MD; Yining Zhao², MD

Corresponding Author:

ShanShan Du, MD Renhe Rehabilitation Hospital Renhe Street Hubei China

Phone: 86 15767853433 Email: <u>1547838432@qq.com</u>

Related Article:

Comment on: http://www.jmir.org/2024/1/e65010/

(J Med Internet Res 2025;27:e69204) doi: 10.2196/69204

KEYWORDS

mobile apps; digital health; atrial fibrillation; anticoagulants; medication adherence; mobile phone

We appreciate the opportunity to read the recently published article by Yoon et al [1], titled Smartphone App for Improving Self-Awareness of Adherence to Edoxaban Treatment in Patients With Atrial Fibrillation (ADHERE-App Trial): Randomized Controlled Trial. The study addresses an important clinical challenge—improving medication adherence in atrial fibrillation patients using digital health interventions. While the findings are valuable, we would like to highlight some additional limitations and considerations that we believe warrant further discussion.

First, the study population was limited to individuals familiar with Android smartphones and the Korean language, likely excluding older adults with lower digital literacy. This presents a significant limitation, as older adults, who often make up a substantial proportion of patients requiring anticoagulation therapy, frequently struggle to adopt and effectively use smartphone-based interventions. For this demographic, navigating apps, managing settings, and understanding instructions on a smartphone can already be challenging, and unfamiliarity with technology exacerbates these difficulties. Such barriers may significantly restrict the practicality and accessibility of these interventions for those who stand to benefit the most. Expanding the target population to include individuals with varying levels of digital literacy and providing tailored support for older adults would greatly enhance the inclusivity and real-world applicability of future research. Additionally, the exclusive focus on edoxaban limits the generalizability of the findings to patients using other anticoagulants with differing dosing regimens and side effects. Broadening the intervention to encompass a more diverse patient population and various

anticoagulants would strengthen the external validity and clinical relevance of the study.

Second, the 6-month follow-up period is another critical consideration, as adherence is a dynamic behavior that often fluctuates over time. While the study demonstrated short-term benefits, long-term follow-up is essential to determine whether these improvements are sustained and lead to meaningful clinical outcomes, such as reduced stroke or bleeding risks [2]. Moreover, relying on pill counts as the primary adherence metric, while practical, does not account for the accuracy of timing or dosage. Using more precise tools, such as electronic medication event monitoring systems, could provide deeper insights into adherence patterns and their clinical implications.

Lastly, although the intervention design is innovative, it could be improved to enhance engagement. Incorporating personalized feedback or adaptive features tailored to individual adherence behaviors might increase user retention and app effectiveness, especially given the higher dropout rates observed in the intervention group. Additionally, integrating the app with complementary strategies, such as caregiver notifications or telemedicine support, could make the intervention more accessible to patients with varying levels of technological proficiency. Addressing these limitations would allow future research to more comprehensively evaluate the app's practicality and clinical impact.

In conclusion, while Yoon et al [1] provide a well-executed study demonstrating the potential of smartphone apps to improve medication adherence, addressing the above considerations could further enhance the clinical relevance and applicability



¹Renhe Rehabilitation Hospital, Hubei, China

²The First Hospital of Shanxi Medical University, Shanxi, China

of their findings. We commend the authors for their research addressing these important aspects. contributions to this evolving field and look forward to future

Conflicts of Interest

None declared.

Editorial Notice

The corresponding author of "Smartphone App for Improving Self-Awareness of Adherence to Edoxaban Treatment in Patients With Atrial Fibrillation (ADHERE-App Trial): Randomized Controlled Trial" declined to respond to this letter.

References

- 1. Yoon M, Lee JH, Kim I, Lee J, Kim M, Kim H, et al. Smartphone App for Improving Self-Awareness of Adherence to Edoxaban Treatment in Patients With Atrial Fibrillation (ADHERE-App Trial): randomized controlled trial. J Med Internet Res. Nov 21, 2024;26:e65010. [FREE Full text] [doi: 10.2196/65010] [Medline: 39570659]
- 2. Hatori Y, Sakai H, Hatori N, Kunishima T, Namiki A, Shimizu M, et al. Long-term outcome and risk factors associated with events in patients with atrial fibrillation treated with oral anticoagulants: The ASSAF-K registry. J Cardiol. Apr 2023;81(4):385-389. [FREE Full text] [doi: 10.1016/j.jjcc.2022.08.012] [Medline: 36109256]

Edited by T Leung; this is a non-peer-reviewed article. Submitted 24.11.24; accepted 31.01.25; published 14.03.25.

Please cite as:

Du S, Zhao Y

Enhancing Digital Health Interventions for Medication Adherence: Considerations for Broader Applicability and Long-Term Impact J Med Internet Res 2025;27:e69204

URL: https://www.jmir.org/2025/1/e69204

doi: 10.2196/69204

PMID:

©ShanShan Du, Yining Zhao. Originally published in the Journal of Medical Internet Research (https://www.jmir.org), 14.03.2025. This is an open-access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research (ISSN 1438-8871), is properly cited. The complete bibliographic information, a link to the original publication on https://www.jmir.org/, as well as this copyright and license information must be included.

